

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
25 August 2005 (25.08.2005)

PCT

(10) International Publication Number
WO 2005/077403 A1

(51) International Patent Classification⁷: **A61K 38/18**

(21) International Application Number:
PCT/US2005/003229

(22) International Filing Date: 4 February 2005 (04.02.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/542,260 4 February 2004 (04.02.2004) US

(71) Applicant (for all designated States except US):
STRYKER CORPORATION [US/US]; 2725 Fair-
field Road, Kalamazoo, Michigan 49005 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **LEE, John C.**
[US/US]; 1119 Haltown Drive, San Antonio, Texas 78213
(US). **YEH, Lee-Chuan C.** [US/US]; 8027 Indian Bend,
San Antonio, Texas 78250 (US).

(74) Agents: **HALEY, James F. Jr.** et al.; FISH & NEAVE
IP GROUP, ROPES & GRAY LLP, 1251 Avenue of the
Americas, New York, New York 10020 (US).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,
MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG,
PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM,
TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM,
ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted
a patent (Rule 4.17(ii)) for the following designations AE,
AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ,
CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE,
EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS,
JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA,
MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM,
PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY,
TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA,
ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ,
NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM,
AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT,
BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR,
HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK,
TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, ML, MR, NE, SN, TD, TG)
- as to the applicant's entitlement to claim the priority of the
earlier application (Rule 4.17(iii)) for all designations

Published:

- with international search report
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: THERAPEUTIC METHODS USING SMADS



at a target locus in a mammal comprising the step of administering to the target locus a Smad are also provided.

(57) Abstract: Methods of inducing the expression of a Smad in a cell or tissue comprising the step of contacting the cell or tissue capable of expressing the Smad with a bone morphogenic protein are provided. Methods of inducing tissue formation and repairing a tissue defect or regenerating tissue,



WO 2005/077403 A1

Therapeutic Methods Using Smads

Technical Field of the Invention

[0001] The present invention relates to methods for tissue formation, repair and regeneration using Smads.

5 Background of the Invention

[0002] The TGF- β superfamily represents a large number of evolutionarily conserved proteins with diverse activities involved in growth, differentiation, cell migration, development, apoptosis and tissue
10 morphogenesis and repair. This large family includes the bone morphogenic proteins (BMPs), TGF- β s and activins. Each subgroup of proteins initiates a unique signaling cascade activated by the formation of a complex with a receptor. The receptors for this family
15 of proteins are type I also known as activin receptor-like kinases (ALKs) and type II serine/threonine kinases. Several such receptors have been identified thus far. The type II receptors are constitutively active. Upon ligand binding the type II receptor
20 phosphorylates particular serine and threonine residues in the type I receptor. The type I serine/threonine

kinases become activated and transduce signals to downstream molecules.

[0003] The downstream molecules in the signaling cascade include the Smads. These molecules are vertebrate counterparts to the *Drosophila* and *Caenorhabditis elegans*, proteins known as Mad (*mothers against dpp*) and Sma, respectively. The name Smad originates from a fusion between Mad and Sma. In recent years, several Smads have been identified (Smad 1-8) (Derynck R. et al., 1996, "Nomenclature: Vertebrate mediators of TGF- β family signals", Cell, 18, 173). The Smads can be divided into three groups: receptor-regulated Smads (R-Smads), common-partner Smads (Co-Smads) and inhibitory Smads (I-Smads). R-Smads are bound to the cell membrane through membrane bound proteins. They transiently interact with and are activated by phosphorylation by activated type I receptor kinases. R-Smads include Smad1, Smad2, Smad3, Smad5 and Smad8. Of those, Smad2 and Smad3 are TGF- β - and activin-specific, whereas Smad1, Smad5 and Smad8 are BMP-specific. The activated R-Smads recruit Co-Smads and these heteromeric complexes translocate to the nucleus. Co-Smads include Smad4. These nuclear Smad complexes bind to DNA directly or indirectly through other DNA-binding proteins, and regulate the transcription of target genes. I-Smads interact with the activated type I receptor, and prevent R-Smads from interacting with activated type I receptors. I-Smads include Smad6 and Smad7.

[0004] As described above, the TGF- β superfamily of proteins have important roles in various physiological events including the inductive properties of the proteins belonging to the BMP family. Therefore, there

remains a need for identifying means useful for promoting tissue regeneration in patients with traumas caused, for example, by injuries or degenerative disorders.

5 **[0005]** The ability to induce Smad protein expression in sufficient quantities at a target locus would be very useful in orthopedic medicine, certain types of plastic surgery, dental and various periodontal and craniofacial reconstructive procedures, and procedures generally
10 involving bone, cartilage, tendon, ligament and neural regeneration. Several Smad genes are now cloned and may be recombinantly expressed in a variety of host systems. The ability to recombinantly produce active Smads, including variants and fragments thereof, and to express
15 them at a target locus makes potential therapeutic treatments using these proteins either alone or together with BMPs feasible.

Summary of the Invention

[0006] This invention is based on the discovery that
20 Smad expression is induced in the presence of various bone morphogenic proteins such as OP-1 (BMP-7) and CDMPs. Therefore, this invention provides a method of inducing the expression of a Smad in a cell or tissue comprising the step of contacting the cell or tissue
25 capable of expressing a Smad with a bone morphogenic protein (BMP). In some embodiments, the tissue is selected from the group consisting of bone, cartilage, tendon, ligament and neural tissue. In one preferred embodiment, the tissue is bone or cartilage. In another
30 preferred embodiment, the tissue is tendon or ligament. In a more preferred embodiment, the tissue is bone. In

another more preferred embodiment, the tissue is cartilage.

[0007] In some embodiments, the cells used in the methods of this invention are progenitor cells. In some
5 embodiments, the progenitor cells include an osteoprogenitor cell, a cartilage progenitor cell, a ligament progenitor cell, a tendon progenitor cell, or a neural progenitor cell. In a preferred embodiment, the progenitor cell is an osteoprogenitor cell or a
10 cartilage progenitor cell. In another preferred embodiment, the progenitor cell is a tendon progenitor cell or a ligament progenitor cell. In a more preferred embodiment, the progenitor cell is an osteoprogenitor cell. In another more preferred embodiment, the
15 progenitor cell is a cartilage progenitor cell.

[0008] In some embodiments, the cell or tissue is contacted with more than one BMP. In some embodiments, the cell or tissue is contacted with two BMPs. BMPs include, but are not limited to, OP-1 (BMP-7), OP-2, OP-
20 3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-
25 12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, and NEURAL. In one preferred embodiment, the BMP is OP-1 (BMP-7). In another preferred embodiment, the BMP is CDMP-1 or GDF-5. In yet another preferred embodiment, the first bone
30 morphogenic protein is OP-1 and the second bone morphogenic protein is CDMP-1 or GDF-5.

[0009] In some embodiments, the Smads used in the methods of the present invention include Smad1, Smad2,

Smad3, Smad5 and Smad8. In a preferred embodiment, the Smad is Smad5. In another preferred embodiment, the Smad is a recombinant Smad.

[0010] In some embodiments, the cell or tissue used
5 in the method of inducing the expression of a Smad in a cell or tissue is further capable of expressing a serine/threonine kinase receptor. In some embodiments the serine/threonine kinase receptor is selected from the group consisting of type I and type II receptors.
10 In some embodiments, only type I receptors are used. In some embodiments, only type II receptors are used. In some embodiments, both type I and type II receptors are used. Preferably, the type I and type II receptors are recombinant. In some embodiments the serine/threonine
15 kinase receptors are linked to an expression control sequence. In some embodiments, the expression control sequence comprises a constitutive promoter. In some embodiments, the expression control sequence comprises an inducible promoter. In some embodiments, the type I
20 receptor is an activin receptor-like kinase (ALK). ALKs include but are not limited to ALK-1, ALK-2, ALK-3, ALK-4, ALK-5, ALK-6, ALK-7 and fragments thereof.

[0011] The invention also provides gene therapy
methods of inducing tissue formation, repairing a tissue
25 defect or regenerating tissue at a target locus. In some embodiments, the invention provides a method of inducing tissue formation at a target locus in a mammal comprising the step of administering to the target locus a nucleic acid encoding a Smad. In some embodiments,
30 the invention provides a method of inducing tissue formation at a target locus in a mammal comprising the step of administering to the target locus a vector comprising a nucleic acid encoding a Smad operably

linked to an expression control sequence. In some
embodiments, the invention provides a method of inducing
tissue formation at a target locus in a mammal
comprising the step of administering to the target locus
5 a cell comprising a vector comprising a nucleic acid
encoding a Smad operably linked to an expression control
sequence.

[0012] The invention further provides a method of
repairing a tissue defect or regenerating tissue at a
10 target locus in a mammal comprising the step of
administering to the target locus a nucleic acid
encoding a Smad. In some embodiments, the invention
provides a method of repairing a tissue defect or
regenerating tissue at a target locus in a mammal
15 comprising the step of administering to the target locus
a vector comprising a nucleic acid encoding a Smad
operably linked to an expression control sequence. In
some embodiments, the invention provides a method of
repairing a tissue defect or regenerating tissue at a
20 target locus in a mammal comprising the step of
administering to the target locus a cell comprising a
vector comprising a nucleic acid encoding a Smad
operably linked to an expression control sequence.

[0013] In some embodiments, the tissue in the gene
25 therapy methods of the present invention is selected
from the group consisting of bone, cartilage, tendon,
ligament and neural tissue. In one preferred
embodiment, the tissue is bone or cartilage. In another
preferred embodiment, the tissue is tendon or ligament.
30 In a more preferred embodiment, the tissue is bone. In
another more preferred embodiment, the tissue is
cartilage.

[0014] In some embodiments, the cells used in the gene therapy methods of this invention are progenitor cells. In some embodiments, the progenitor cells include an osteoprogenitor cell, a cartilage progenitor cell, a ligament progenitor cell, a tendon progenitor cell, or a neural progenitor cell. In a preferred embodiment, the progenitor cell is an osteoprogenitor cell or a cartilage progenitor cell. In another preferred embodiment, the progenitor cell is a tendon progenitor cell or a ligament progenitor cell. In a more preferred embodiment, the progenitor cell is an osteoprogenitor cell. In another more preferred embodiment, the progenitor cell is a cartilage progenitor cell.

[0015] In some embodiments, the expression control sequence operably linked to the nucleic acid encoding a Smad comprises a constitutive promoter. In other embodiments, the expression control sequence operably linked to the nucleic acid encoding a Smad comprises an inducible promoter. A Smad according to this invention includes, but is not limited to, Smad1, Smad2, Smad3, Smad5, Smad8 and fragments thereof. In a preferred embodiment, the Smad is Smad5. In another preferred embodiment, the Smad is a recombinant Smad.

[0016] In some embodiments, the methods of inducing tissue formation, repairing a tissue defect or regeneration tissue of this invention further comprise the step of administering to the target locus a serine/threonine kinase receptor. In some embodiments, a nucleic acid encoding a serine/threonine kinase receptor is administered. In some embodiments, a vector comprising a nucleic acid encoding a serine/threonine kinase receptor operably linked to an expression control

sequence is administered. In other embodiments, a cell comprising a vector comprising a nucleic acid encoding a serine/threonine kinase receptor operably linked to an expression control sequence is administered.

5 [0017] In some embodiments, the expression control sequence operably linked to the a serine/threonine kinase receptor comprises a constitutive promoter. In some embodiments, the expression control sequence operably linked to the serine/threonine kinase receptor
10 comprises an inducible promoter.

 [0018] In some embodiments, the serine/threonine kinase receptor used in the gene therapy methods of inducing tissue formation, repairing a tissue defect or regenerating tissue at a target locus, is selected from
15 the group consisting of type I and type II receptors. In some embodiments, only type I receptors are used. In some embodiments, only type II receptors are used. In some embodiments both type I and type II receptors are used. Preferably recombinant type I and recombinant
20 type II receptors are used. In some embodiments, the type I receptor is an activin receptor-like kinase (ALK). The ALKs that may be used in the present invention include, but are not limited to ALK-1, ALK-2, ALK-3, ALK-4, ALK-5, ALK-6, ALK-7, and fragments
25 thereof. In a preferred embodiment, the ALK is ALK-2, ALK-3, ALK-6 and fragments thereof.

 [0019] In some embodiments, the methods of inducing tissue formation, repairing a tissue defect or regeneration tissue of this invention further comprise
30 the step of administering to the target locus a bone morphogenic protein. In some embodiments, the bone morphogenic protein is administered as a nucleic acid. In some embodiments, the bone morphogenic protein is

administered as a vector comprising a nucleic acid encoding the bone morphogenic protein operably linked to an expression control sequence. In some embodiments, the bone morphogenic protein is administered as a cell
5 comprising a vector comprising a nucleic acid encoding a bone morphogenic protein operably linked to an expression control sequence. In some embodiments, the expression control sequence linked to the BMP nucleic acid comprises a constitutive promoter. In some
10 embodiments, the expression control sequence linked to the BMP nucleic acid comprises an inducible promoter.

[0020] The bone morphogenic protein according to this invention includes, but is not limited to, OP-1 (BMP-7), OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-
15 BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A
20 protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, and fragments thereof. In some embodiments, one BMP is administered. In some embodiments, more than one BMP is administered. In some embodiments, two BMPs are administered. In some embodiments, the first BMP is OP-
25 1 and the second BMP is CDMP-1 or GDF-5. In a preferred embodiment, the BMP is OP-1. In another preferred embodiment, the BMP is CDMP-1 or GDF-5.

Brief Description of the Drawings

[0021] Figure 1 is a Western blot analysis for Smad5
30 protein in C2C12 cells following treatment with OP-1 and CDMP-1. Lane 1 is the Smad5 levels in control C2C12 cells. Lane 2 is the Smad5 levels in CDMP-1 treated

(200 ng/ml) C2C12 cells. Lane 3 is the Smad5 levels in OP-1 treated (100 ng/ml) C2C12 cells. Lane 4 is the Smad5 levels in cells treated with both OP-1 (100 ng/ml) and CDMP-1 (200 ng/ml).

5 [0022] Figure 2 is a plasmid map of pW24 containing the OP-1 gene.

Detailed Description of the Invention

[0023] In order that the invention herein described may be fully understood, the following detailed
10 description is set forth.

[0024] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as those commonly understood by one of ordinary skill in the art to which this invention belongs. Although
15 methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. The materials, methods and examples are illustrative only, and are not intended to
20 be limiting. All publications, patents and other documents mentioned herein are incorporated by reference in their entirety.

[0025] Throughout this specification, the word "comprise" or variations such as "comprises" or
25 "comprising" will be understood to imply the inclusion of a stated integer or groups of integers but not the exclusion of any other integer or group of integers.

[0026] In order to further define the invention, the following terms and definitions are provided herein.

30 [0027] The term "defect" or "defect site," refers to a disruption of the specified tissue. A defect can assume the configuration of a "void", which is

understood to mean a three-dimensional defect such as, for example, a gap, cavity, hole or other substantial disruption in the structural integrity of the tissue (e.g., bone, chondral, osteochondral, neural, ligament, tendon). Moreover, a defect can also be a detachment of the tendon or ligament from its point of attachment to bone, cartilage or muscle. In certain embodiments, the defect is such that it is incapable of endogenous or spontaneous repair. A defect can be the result of accident, disease, and/or surgical manipulation.

[0028] The term "target locus" refers to the site (e.g., a defect) in any tissue that is in need of repair or regeneration. The target locus need not be a defect site. It may be any site where bone, cartilage, tendon, ligament or neural tissue regeneration is desired.

[0029] The term "repair" refers to new tissue formation which is sufficient to at least partially fill the void or structural discontinuity at the defect site. Repair does not, however, mean, or otherwise necessitate, a process of complete healing or a treatment, which is 100% effective at restoring a defect to its pre-defect physiological/structural/mechanical state.

[0030] The term "therapeutically effective amount" refers to an amount effective to repair, regenerate, promote, accelerate, prevent degradation, or form tissue.

[0031] The term "patient" refers to an animal, including a mammal (e.g., a human).

[0032] The term "bone morphogenic protein (BMP)" refers to a protein belonging to the BMP family of the TGF- β superfamily of proteins (BMP family) based on DNA and amino acid sequence homology. A protein belongs to

the BMP family according to this invention when it has at least 50% amino acid sequence identity with at least one known BMP family member within the conserved C-terminal cysteine-rich domain, which characterizes the BMP protein family. Preferably, the protein has at least 70% amino acid sequence identity with at least one known BMP family member within the conserved C-terminal cysteine rich domain. Members of the BMP family may have less than 50% DNA or amino acid sequence identity overall. Bone morphogenic proteins may be monomeric, homo- or hetero-dimeric. Bone morphogenic proteins include osteogenic proteins.

[0033] Bone morphogenic proteins are capable of inducing progenitor cells to proliferate and/or to initiate differentiation pathways that lead to cartilage, bone, tendon, ligament or other types of tissue formation depending on local environmental cues, and thus bone morphogenic proteins may behave differently in different surroundings. For example, a bone morphogenic protein may induce bone tissue at one treatment site and cartilage tissue at a different treatment site. Bone morphogenic proteins include full length proteins as well as fragments thereof.

[0034] The term "osteogenic protein (OP)" refers to a bone morphogenic protein that is capable of inducing a progenitor cell to form cartilage and/or bone. The bone may be intramembranous bone or endochondral bone. Most osteogenic proteins are members of the BMP protein family and are thus also BMPs. As described elsewhere herein, the class of proteins is typified by human osteogenic protein (hOP-1). Other osteogenic proteins useful in the practice of the invention include but are not limited to, osteogenically active forms of OP-1, OP-

2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL, conservative amino acid sequence variants thereof having osteogenic activity and fragments thereof. In one currently preferred embodiment, osteogenic protein includes any one of: OP-1, CDMP-1, CDMP-2, CDMP-3, GDF-5, GDF-6, GDF-7, amino acid sequence variants and homologs thereof, including species homologs thereof and fragments thereof. Particularly preferred osteogenic proteins are those comprising an amino acid sequence having at least 70% homology with the C-terminal 102-106 amino acids, defining the conserved seven cysteine domain, of, e.g., human OP-1. Certain preferred embodiments of the instant invention comprise the osteogenic protein, OP-1. As further described elsewhere herein, the osteogenic proteins suitable for use with this invention can be identified by means of routine experimentation using the art-recognized bioassay described by Reddi and Sampath (Sampath et al., *Proc. Natl. Acad. Sci.*, 84, pp. 7109-13, incorporated herein by reference).

[0035] Proteins useful in this invention include eukaryotic proteins identified as osteogenic proteins (see U.S. Patent 5,011,691, incorporated herein by reference), such as the OP-1, OP-2, OP-3 and CBMP-2 proteins, as well as amino acid sequence-related proteins, such as DPP (from *Drosophila*), Vg1 (from *Xenopus*), Vgr-1 (from mouse), GDF-1 (from humans, see

Lee, *PNAS*, 88, pp. 4250-4254 (1991)), 60A (from
Drosophila, see Wharton et al., *PNAS*, 88, pp. 9214-9218
(1991)), dorsalin-1 (from chick, see Basler et al.,
Cell, 73, pp. 687-702 (1993) and GenBank accession
5 number L12032) and GDF-5 (from mouse, see Storm et al.,
Nature, 368, pp. 639-643 (1994)). The teachings of the
above references are incorporated herein by reference.
BMP-3 is also preferred. Additional useful proteins
include biosynthetic morphogenic constructs disclosed in
10 U.S. Pat. No. 5,011,691, incorporated herein by
reference, e.g., COP-1, COP-3, COP-4, COP-5, COP-7 and
COP-16, as well as other proteins known in the art.
Still other proteins include osteogenically active forms
of BMP-3b (see Takao, et al., *Biochem. Biophys. Res.*
15 *Comm.*, 219, pp. 656-662 (1996)). BMP-9 (see WO
95/33830), BMP-15 (see WO 96/35710), BMP-12 (see WO
95/16035), CDMP-1 (see WO 94/12814), CDMP-2 (see WO
94/12814), BMP-10 (see WO 94/26893), GDF-1 (see WO
92/00382), GDF-10 (see WO95/10539), GDF-3 (see WO
20 94/15965) and GDF-7 (see WO95/01802). The teachings of
the above references are incorporated herein by
reference. BMPs (identified by sequence homology) must
have demonstrable osteogenic activity in a functional
bioassay to be osteogenic proteins according to this
25 invention.

[0036] The term "amino acid sequence homology" is
understood to include both amino acid sequence identity
and similarity. Homologous sequences share identical
and/or similar amino acid residues, where similar
30 residues are conservative substitutions for, or "allowed
point mutations" of, corresponding amino acid residues
in an aligned reference sequence. Thus, a candidate
polypeptide sequence that shares 70% amino acid homology

with a reference sequence is one in which any 70% of the aligned residues are either identical to, or are conservative substitutions of, the corresponding residues in a reference sequence. Certain particularly preferred bone morphogenic polypeptides share at least 60%, and preferably 70% amino acid sequence identity with the C-terminal 102-106 amino acids, defining the conserved seven-cysteine domain of human OP-1 and related proteins.

10 **[0037]** Amino acid sequence homology can be determined by methods well known in the art. For instance, to determine the percent homology of a candidate amino acid sequence to the sequence of the seven-cysteine domain, the two sequences are first aligned. The alignment can
15 be made with, e.g., the dynamic programming algorithm described in Needleman et al., J. Mol. Biol., 48, pp. 443 (1970), and the Align Program, a commercial software package produced by DNASTar, Inc. The teachings by both sources are incorporated by reference
20 herein. An initial alignment can be refined by comparison to a multi-sequence alignment of a family of related proteins. Once the alignment is made and refined, a percent homology score is calculated. The aligned amino acid residues of the two sequences are
25 compared sequentially for their similarity to each other. Similarity factors include similar size, shape and electrical charge. One particularly preferred method of determining amino acid similarities is the PAM250 matrix described in Dayhoff et al., Atlas of
30 Protein Sequence and Structure, 5, pp. 345-352 (1978 & Supp.), which is incorporated herein by reference. A similarity score is first calculated as the sum of the aligned pair wise amino acid similarity scores.

Insertions and deletions are ignored for the purposes of percent homology and identity. Accordingly, gap penalties are not used in this calculation. The raw score is then normalized by dividing it by the geometric mean of the scores of the candidate sequence and the seven-cysteine domain. The geometric mean is the square root of the product of these scores. The normalized raw score is the percent homology.

[0038] The term "conservative substitutions" refers to residues that are physically or functionally similar to the corresponding reference residues. That is, a conservative substitution and its reference residue have similar size, shape, electric charge, chemical properties including the ability to form covalent or hydrogen bonds, or the like. Preferred conservative substitutions are those fulfilling the criteria defined for an accepted point mutation in Dayhoff *et al.*, *supra*. Examples of conservative substitutions are substitutions within the following groups: (a) valine, glycine; (b) glycine, alanine; (c) valine, isoleucine, leucine; (d) aspartic acid, glutamic acid; (e) asparagine, glutamine; (f) serine, threonine; (g) lysine, arginine, methionine; and (h) phenylalanine, tyrosine. The term "conservative variant" or "conservative variation" also includes the use of a substituting amino acid residue in place of an amino acid residue in a given parent amino acid sequence, where antibodies specific for the parent sequence are also specific for, i.e., "cross-react" or "immuno-react" with, the resulting substituted polypeptide sequence.

[0039] The term "fragment thereof" or "fragment" refers to a stretch of at least about 5 amino acid residues. In some embodiments, this term refers to a

stretch of at least about 10 amino acid residues. In other embodiments, it refers to a stretch of at least about 15 to 20 amino acid residues. The fragments may be naturally derived or synthetically generated. To be
5 active, any fragment must have sufficient length to display biological activity.

Methods Using Smads

[0040] The present invention provides a method of inducing the expression of a Smad in a cell or tissue
10 comprising the step of contacting the cell or tissue capable of expressing the Smad with a bone morphogenic protein.

[0041] This invention also provides gene therapy methods for inducing tissue formation at a target locus,
15 repairing a tissue defect or regenerating tissue at a target locus using Smads. In some embodiments, the methods comprise the step of administering to the target locus a nucleic acid encoding a Smad. In some embodiments, the methods comprise the step of
20 administering to the target locus a vector comprising a nucleic acid encoding a Smad operably linked to an expression control sequence. In some embodiments, the methods comprise the step of administering a cell comprising a vector comprising a nucleic acid encoding a
25 Smad operably linked to an expression control sequence. In some embodiments, the expression control sequence operably linked to a Smad nucleic acid comprises an inducible promoter. In some embodiments, the expression control sequence operably linked to a Smad nucleic acid
30 comprises a constitutive promoter.

[0042] A Smad according to the present invention is a R-Smad. R-Smads includes, but are not limited to,

Smad1, Smad2, Smad3, Smad5, Smad8 or fragments thereof. In a preferred embodiment, the Smad is Smad5. In another embodiment the Smad is recombinant.

[0043] In some embodiments, the gene therapy methods further comprise the step of administering to the target locus a pharmaceutically effective amount of a BMP. In some embodiments, the methods further comprise the step of administering a nucleic acid encoding a BMP. In some embodiments, the methods further comprise the step of administering a vector comprising a nucleic acid encoding a BMP operably linked to an expression control sequence. In some embodiments, the methods further comprise the step of administering a cell comprising a vector comprising a nucleic acid encoding a BMP operably linked to an expression control sequence. In some embodiments, the expression control sequence operably linked to a BMP nucleic acid comprises an inducible promoter. In some embodiments, the expression control sequence operably linked to a BMP nucleic acid comprises a constitutive promoter.

[0044] The BMPs according to the invention include but are not limited to, OP-1 (BMP-7), OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, and NEURAL. (see *infra*, for discussion of BMPs). In a preferred embodiment, the BMP is OP-1 (BMP-7). In another preferred embodiment, the BMP is CDMP-1 or GDF-5.

[0045] In some embodiments, more than one BMP is administered. In one preferred embodiment, two BMPs are administered. In yet another preferred embodiment, the first BMP is OP-1 and the second BMP is CDMP-1 or GDF-5.
5 In another embodiment three BMPs are administered. In some embodiments, the BMP is recombinant.

[0046] In some embodiments, the gene therapy methods of this invention further comprise the step of administering a serine/threonine kinase receptor. In
10 some embodiments, the methods of this invention further comprise the step of administering to the target locus a nucleic acid encoding a serine/threonine kinase receptor. In some embodiments, the methods of this invention further comprise the step of administering to
15 the target locus a vector comprising a nucleic acid encoding a serine/threonine kinase receptor operably linked to an expression control sequence. In some embodiments, the methods of this invention further comprise the step of administering to the target locus a cell comprising a vector comprising a nucleic acid
20 encoding a serine/threonine kinase receptor operably linked to an expression control sequence. In some embodiments, the serine/threonine kinase receptor is selected from the group consisting of type I and type II receptors. In some embodiments, only a type I receptor
25 is used. In some embodiments, only a type II receptors is used. In some embodiments both type I and type II receptors are used. Preferably the type I and type II receptors are recombinant. In some embodiments, the
30 expression control sequence operably linked to a serine/threonine kinase receptor nucleic acid comprises an inducible promoter. In some embodiments, the expression control sequence operably linked to a

serine/threonine kinase receptor nucleic acid comprises a constitutive promoter.

[0047] In some embodiments, the type I receptor is an activin receptor-like kinase (ALK). The ALKs according to this invention include, but are not limited to ALK-1, 5 ALK-2, ALK-3, ALK-4, ALK-5, ALK-6, ALK-7, and fragments thereof. Preferred ALKs are ALK-2, ALK-3 and ALK-6.

Bone Morphogenic Protein (BMP) Family

[0048] The BMP family, named for its representative 10 bone morphogenic/osteogenic protein family members, belongs to the TGF- β protein superfamily. Of the reported "BMPs" (BMP-1 to BMP-18), isolated primarily based on sequence homology, all but BMP-1 remain classified as members of the BMP family of morphogenic 15 proteins (Ozkaynak et al., *EMBO J.*, 9, pp. 2085-93 (1990)).

[0049] The BMP family includes other structurally-related members which are bone morphogenic proteins, including the *drosophila* decapentaplegic gene complex 20 (DPP) products, the Vg1 product of *Xenopus laevis* and its murine homolog, Vgr-1 (see, e.g., Massagué, *Annu. Rev. Cell Biol.*, 6, pp. 597-641 (1990), incorporated herein by reference).

[0050] The *Drosophila* DPP and *Xenopus* Vg-1 gene 25 products are 50% identical to each other (and 35-40% identical to TGF- β). Both the Dpp and Vg-1 products are morphogenic proteins that participate in early patterning events during embryogenesis of their respective hosts. These products appear to be most 30 closely related to mammalian bone morphogenetic proteins BMP-2 and BMP-4, whose C-terminal domains are 75% identical with that of Dpp.

[0051] The C-terminal domains of BMP-3, BMP-5, BMP-6, and OP-1 (BMP-7) are about 60% identical to that of BMP-2, and the C-terminal domains of BMP-6 and OP-1 are 87% identical. BMP-6 is likely the human homolog of the murine Vgr-1 (Lyons *et al.*, *Proc. Natl. Acad. Sci. U.S.A.*, 86, pp. 4554-59 (1989)); the two proteins are 92% identical overall at the amino acid sequence level (U.S. Patent No. 5,459,047, incorporated herein by reference). BMP-6 is 58% identical to the *Xenopus* Vg-1 product.

[0052] The naturally occurring bone morphogenic proteins share substantial amino acid sequence homology in their C-terminal regions (domains). Typically, the above-mentioned naturally occurring osteogenic proteins are translated as a precursor, having an N-terminal signal peptide sequence typically less than about 30 residues, followed by a "pro" domain that is cleaved to yield the mature C-terminal domain of approximately 100-140 amino acids. The signal peptide is cleaved rapidly upon translation, at a cleavage site that can be predicted in a given sequence using the method of Von Heijne, *Nucleic Acids Research*, 14, pp. 4683-4691 (1986). The pro domain typically is about three times larger than the fully processed mature C-terminal domain.

[0053] Another characteristic of the BMP protein family members is their ability to dimerize. Several bone-derived osteogenic proteins (OPs) and BMPs are found as homo- and heterodimers in their active forms. The ability of OPs and BMPs to form heterodimers may confer additional or altered morphogenic inductive capabilities on bone morphogenic proteins. Heterodimers may exhibit qualitatively or quantitatively different

binding affinities than homodimers for OP and BMP receptor molecules. Altered binding affinities may in turn lead to differential activation of receptors that mediate different signaling pathways, which may ultimately lead to different biological activities or outcomes. Altered binding affinities could also be manifested in a tissue or cell type-specific manner, thereby inducing only particular progenitor cell types to undergo proliferation and/or differentiation.

10 **[0054]** In one preferred embodiment of this invention, the BMPs independently comprise a pair of subunits disulfide bonded to produce a dimeric species, wherein at least one of the subunits comprises a recombinant peptide belonging to the BMP protein family. In another preferred embodiment of this invention, the BMPs independently comprise a pair of subunits that produce a dimeric species formed through non-covalent interactions, wherein at least one of the subunits comprises a recombinant peptide belonging to the BMP protein family. Non-covalent interactions include Van der Waals, hydrogen bond, hydrophobic and electrostatic interactions. The dimeric species may be a homodimer or heterodimer and is capable of inducing cell proliferation and/or tissue formation. In some embodiments, the BMPs are each independently monomers.

25 **[0055]** In preferred embodiments, the pair of morphogenic polypeptides have amino acid sequences each comprising a sequence that shares a defined relationship with an amino acid sequence of a reference morphogen. Herein, preferred osteogenic polypeptides share a defined relationship with a sequence present in osteogenically active human OP-1, SEQ ID NO: 1. However, any one or more of the naturally occurring or

biosynthetic sequences disclosed herein similarly could be used as a reference sequence. Preferred osteogenic polypeptides share a defined relationship with at least the C-terminal six cysteine domain of human OP-1, residues 335-431 of SEQ ID NO: 1. Preferably, osteogenic polypeptides share a defined relationship with at least the C-terminal seven cysteine domain of human OP-1, residues 330-431 of SEQ ID NO: 1. That is, preferred polypeptides in a dimeric protein with bone morphogenic activity each comprise a sequence that corresponds to a reference sequence or is functionally equivalent thereto.

[0056] Functionally equivalent sequences include functionally equivalent arrangements of cysteine residues disposed within the reference sequence, including amino acid insertions or deletions which alter the linear arrangement of these cysteines, but do not materially impair their relationship in the folded structure of the dimeric morphogen protein, including their ability to form such intra- or inter-chain disulfide bonds as may be necessary for morphogenic activity. Functionally equivalent sequences further include those wherein one or more amino acid residues differs from the corresponding residue of a reference sequence, e.g., the C-terminal seven cysteine domain (also referred to herein as the conserved seven cysteine skeleton) of human OP-1, provided that this difference does not destroy bone morphogenic activity. Accordingly, conservative substitutions of corresponding amino acids in the reference sequence are preferred. Particularly preferred conservative substitutions are those fulfilling the criteria defined for an accepted

point mutation in Dayhoff *et al.*, *supra*, the teachings of which are incorporated by reference herein.

[0057] The osteogenic protein OP-1 has been described (see, *e.g.*, Oppermann *et al.*, U. S. Patent No. 5,354,557, incorporated herein by reference). Natural-sourced osteogenic protein in its mature, native form is a glycosylated dimer typically having an apparent molecular weight of about 30-36 kDa as determined by SDS-PAGE. When reduced, the 30 kDa protein gives rise to two glycosylated peptide subunits having apparent molecular weights of about 16 kDa and 18 kDa. The unglycosylated protein, which also has osteogenic activity, has an apparent molecular weight of about 27 kDa. When reduced, the 27 kDa protein gives rise to two unglycosylated polypeptides, having molecular weights of about 14 kDa to 16 kDa, capable of inducing endochondral bone formation in a mammal. Osteogenic proteins may include forms having varying glycosylation patterns, varying N-termini, and active truncated or mutated forms of native protein.

[0058] As described above, particularly useful sequences include those comprising the C-terminal 96 or 102 amino acid sequences of DPP (from *Drosophila*), Vg1 (from *Xenopus*), Vgr-1 (from mouse), the OP-1 and OP-2 proteins, (see U.S. Pat. No. 5,011,691 and Oppermann *et al.*, incorporated herein by reference), as well as the proteins referred to as BMP-2, BMP-3, BMP-4 (see WO 88/00205, U.S. Patent No. 5,013,649 and WO 91/18098, incorporated herein by reference), BMP-5 and BMP-6 (see WO 90/11366, PCT/US90/01630, incorporated herein by reference), BMP-8 and BMP-9.

[0059] Preferred BMPs of this invention comprise at least one polypeptide selected from the group consisting

of OP-1 (BMP-7), OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, 5 GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, NEURAL and amino acid sequence variants and homologs thereof, including species homologs thereof and fragments 10 thereof. In some embodiments, one BMP is used. In some embodiments, more than one BMP is used. In some embodiments, two BMPs are used. In some embodiments, three BMPs are used. In some embodiments, the first BMP is OP-1 (BMP-7) or a fragment thereof, and the second 15 BMP is selected from the group consisting of CDMP-1, CDMP-2, CDMP-3, GDF-5, GDF-6, GDF-7 and fragments thereof. In some embodiments, the second BMP is CDMP-1, GDF-5 or a fragment thereof. In some embodiments, the first BMP is OP-1 and the second BMP is CDMP-2 or GDF-5.

20 **[0060]** Publications disclosing these sequences, as well as their chemical and physical properties, include: OP-1 and OP-2 (U.S. Patent No. 5,011,691; U.S. Patent No. 5,266,683; Ozkaynak et al., *EMBO J.*, 9, pp. 2085-2093 (1990); OP-3 (WO 94/10203 (PCT US93/10520)), 25 BMP-2, BMP-3, BMP-4, (WO 88/00205; Wozney et al. *Science*, 242, pp. 1528-1534 (1988)), BMP-5 and BMP-6, (Celeste et al., *PNAS*, 87, 9843-9847 (1991)), Vgr-1 (Lyons et al., *PNAS*, 86, pp. 4554-4558 (1989)); DPP (Padgett et al. *Nature*, 325, pp. 81-84 (1987)); Vg-1 30 (Weeks, *Cell*, 51, pp. 861-867 (1987)); BMP-9 (WO95/33830 (PCT/US95/07084); BMP-10 (WO 94/26893 (PCT/US94/05290); BMP-11 (WO 94/26892 (PCT/US94/05288); BMP-12 (WO95/16035 (PCT/US94/14030); BMP-13 (WO95/16035 (PCT/US94/14030);

GDF-1 (WO 92/00382 (PCT/US91/04096) and Lee et al. *PNAS*, 88, pp. 4250-4254 (1991); GDF-8 (WO 94/21681 (PCT/US94/03019); GDF-9 (WO 94/15966 (PCT/US94/00685); GDF-10 (WO 95/10539 (PCT/US94/11440); GDF-11 (WO 5 96/01845 (PCT/US95/08543); BMP-15 (WO 96/36710 (PCT/US96/06540); MP-121 (WO 96/01316 (PCT/EP95/02552); GDF-5 (CDMP-1, MP52) (WO 94/15949 (PCT/US94/00657) and WO 96/14335 (PCT/US94/12814) and WO 93/16099 (PCT/EP93/00350)); GDF-6 (CDMP-2, BMP13) (WO 95/01801 10 (PCT/US94/07762) and WO 96/14335 and WO 95/10635 (PCT/US94/14030)); GDF-7 (CDMP-3, BMP12) (WO 95/10802 (PCT/US94/07799) and WO 95/10635 (PCT/US94/14030)). The above publications are incorporated herein by reference.

[0061] In another embodiment of this invention, the 15 BMPs may be prepared synthetically. BMPs prepared synthetically may be native, or may be non-native proteins, i.e., those not otherwise found in nature. Non-native osteogenic proteins have been synthesized using a series of consensus DNA sequences (U.S. Patent 20 No. 5,324,819, incorporated herein by reference). These consensus sequences were designed based on partial amino acid sequence data obtained from natural osteogenic products and on their observed homologies with other genes reported in the literature having a presumed or 25 demonstrated developmental function.

[0062] Several of the biosynthetic consensus sequences (called consensus osteogenic proteins or "COPs") have been expressed as fusion proteins in prokaryotes. Purified fusion proteins may be cleaved, 30 refolded, implanted in an established animal model and shown to have bone- and/or cartilage-inducing activity. The currently preferred synthetic osteogenic proteins comprise two synthetic amino acid sequences designated ,

COP-5 (SEQ. ID NO: 2) and COP-7 (SEQ. ID NO: 3).

Oppermann et al., U. S. Patent Nos. 5,011,691 and 5,324,819, which are incorporated herein by reference, describe the amino acid sequences of COP-5 and COP-7 as shown below:

COP5 LYVDFS-DVGWDDWIVAPPGYQAFYCHGECPPFLAD

COP7 LYVDFS-DVGWNDWIVAPPGYHAFYCHGECPPFLAD

COP5 HFNSTN--H-AVVQTLVNSVNSKI--PKACCVPTELSA

10 COP7 HLNSTN--H-AVVQTLVNSVNSKI--PKACCVPTELSA

COP5 ISMLYLDENEKVVVKYNQEMVVEGCGCR

COP7 ISMLYLDENEKVVVKYNQEMVVEGCGCR

15 **[0063]** In these amino acid sequences, the dashes (-) are used as fillers only to line up comparable sequences in related proteins. Differences between the aligned amino acid sequences are highlighted.

[0064] The DNA and amino acid sequences of these and
20 other BMP family members are published and may be used by those of skill in the art to determine whether a newly identified protein belongs to the BMP family.

[0065] In certain preferred embodiments, the BMPs useful herein independently include those in which the
25 amino acid sequences comprise a sequence sharing at least 70% amino acid sequence homology or "similarity", preferably 80%, more preferably 90%, even more preferably 95%, even more preferably 98% homology or similarity, with a reference bone morphogenic protein
30 selected from the foregoing naturally occurring proteins. Preferably, the reference protein is human OP-1, and the reference sequence thereof is the C-terminal seven cysteine domain present in osteogenically

active forms of human OP-1, residues 330-431 of SEQ ID NO: 1. In some embodiments, the BMP comprises a dimeric protein having an amino acid sequence having at least 70% homology within the C-terminal 102-106 amino acids of human OP-1. In certain embodiments, a polypeptide suspected of being functionally equivalent to a reference BMP polypeptide is aligned therewith using the method of Needleman, *et al.*, *supra*, implemented conveniently by computer programs such as the Align program (DNASTar, Inc.). As noted above, internal gaps and amino acid insertions in the candidate sequence are ignored for purposes of calculating the defined relationship, conventionally expressed as a level of amino acid sequence homology or identity, between the candidate and reference sequences. In one preferred embodiment, the reference sequence is OP-1. In another preferred embodiment, the reference sequence is selected from CDMP-1, CDMP-2 or CDMP-3. Bone morphogenic proteins useful herein accordingly include allelic, phylogenetic counterpart and other variants of the preferred reference sequence, whether naturally-occurring or biosynthetically produced (e.g., including "muteins" or "mutant proteins"), as well as novel members of the general morphogenic family of proteins, including those set forth and identified above. Certain particularly preferred bone morphogenic polypeptides share at least 60% amino acid identity with the preferred reference sequence of human OP-1, still more preferably at least 65% amino acid identity therewith.

[0066] In another embodiment, useful BMPs include those sharing the conserved seven cysteine domain and sharing at least 70% amino acid sequence homology (similarity) within the C-terminal active domain, as

defined herein. In still another embodiment, the BMPs of the invention can be defined as osteogenically active proteins having any one of the generic sequences defined herein, including OPX (SEQ ID NO: 4) and Generic Sequences 7 (SEQ ID NO: 5) and 8 (SEQ ID NO: 6), or Generic Sequences 9 (SEQ ID NO: 7) and 10 (SEQ ID NO: 8).

[0067] The family of bone morphogenic polypeptides useful in the present invention, and members thereof, can be defined by a generic amino acid sequence. For example, Generic Sequence 7 (SEQ ID NO: 5) and Generic Sequence 8 (SEQ ID NO: 6) are 97 and 102 amino acid sequences, respectively, and accommodate the homologies shared among preferred protein family members identified to date, including at least OP-1, OP-2, OP-3, CBMP-2A, CBMP-2B, BMP-3, 60A, DPP, Vg1, BMP-5, BMP-6, Vgr-1, and GDF-1. The amino acid sequences for these proteins are described herein and/or in the art, as summarized above. The generic sequences include both the amino acid identity shared by these sequences in the C-terminal domain, defined by the six and seven cysteine skeletons (Generic Sequences 7 and 8, respectively), as well as alternative residues for the variable positions within the sequence. The generic sequences provide an appropriate cysteine skeleton where inter- or intramolecular disulfide bonds can form, and contain certain critical amino acids likely to influence the tertiary structure of the folded proteins. In addition, the generic sequences allow for an additional cysteine at position 36 (Generic Sequence 7) or position 41 (Generic Sequence 8), thereby encompassing the morphogenically active sequences of OP-2 and OP-3.

Generic Sequence 7

			Leu	Xaa	Xaa	Xaa	Phe	Xaa	Xaa
			1				5		
Xaa	Gly	Trp	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Pro
		10					15		
Xaa	Xaa	Xaa	Xaa	Ala	Xaa	Tyr	Cys	Xaa	Gly
		20					25		
Xaa	Cys	Xaa	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa
		30					35		
Xaa	Xaa	Xaa	Asn	His	Ala	Xaa	Xaa	Xaa	Xaa
		40					45		
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
		50					55		
Xaa	Xaa	Xaa	Cys	Cys	Xaa	Pro	Xaa	Xaa	Xaa
		60					65		
Xaa	Xaa	Xaa	Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa
		70					75		
Xaa	Xaa	Xaa	Val	Xaa	Leu	Xaa	Xaa	Xaa	Xaa
		80					85		
Xaa	Met	Xaa	Val	Xaa	Xaa	Cys	Xaa	Cys	Xaa
		90					95		

wherein each Xaa independently is selected from a group
 5 of one or more specified amino acids defined as follows:
 "res." means "residue" and Xaa at res.2 = (Tyr or Lys);
 Xaa at res.3 = Val or Ile); Xaa at res.4 = (Ser, Asp or
 Glu); Xaa at res.6 = (Arg, Gln, Ser, Lys or Ala); Xaa at
 res.7 = (Asp or Glu); Xaa at res.8 = (Leu, Val or Ile);
 10 Xaa at res. 11 = (Gln, Leu, Asp, His, Asn or Ser); Xaa
 at res.12 = (Asp, Arg, Asn or Glu); Xaa at res.13 = (Trp
 or Ser); Xaa at res.14 = (Ile or Val); Xaa at res.15 =
 (Ile or Val); Xaa at res.16 (Ala or Ser); Xaa at res.18
 = (Glu, Gln, Leu, Lys, Pro or Arg); Xaa at res.19 = (Gly
 15 or Ser); Xaa at res.20 = (Tyr or Phe); Xaa at res.21 =
 (Ala, Ser, Asp, Met, His, Gln, Leu or Gly); Xaa at
 res.23 = (Tyr, Asn or Phe); Xaa at res.26 = (Glu, His,
 Tyr, Asp, Gln, Ala or Ser); Xaa at res.28 = (Glu, Lys,
 Asp, Gln or Ala); Xaa at res.30 = (Ala, Ser, Pro, Gln,

Ile or Asn); Xaa at res.31 = (Phe, Leu or Tyr); Xaa at
res.33 = (Leu, Val or Met); Xaa at res.34 = (Asn, Asp,
Ala, Thr or Pro); Xaa at res.35 = (Ser, Asp, Glu, Leu,
Ala or Lys); Xaa at res.36 = (Tyr, Cys, His, Ser or
5 Ile); Xaa at res.37 = (Met, Phe, Gly or Leu); Xaa at
res.38 = (Asn, Ser or Lys); Xaa at res.39 = (Ala, Ser,
Gly or Pro); Xaa at res.40 = (Thr, Leu or Ser); Xaa at
res.44 = (Ile, Val or Thr); Xaa at res.45 = (Val, Leu,
Met or Ile); Xaa at res.46 = (Gln or Arg); Xaa at res.47
10 = (Thr, Ala or Ser); Xaa at res.48 = (Leu or Ile); Xaa
at res.49 = (Val or Met); Xaa at res.50 = (His, Asn or
Arg); Xaa at res.51 = (Phe, Leu, Asn, Ser, Ala or Val);
Xaa at res.52 = (Ile, Met, Asn, Ala, Val, Gly or Leu);
Xaa at res.53 = (Asn, Lys, Ala, Glu, Gly or Phe); Xaa at
15 res.54 = (Pro, Ser or Val); Xaa at res.55 = (Glu, Asp,
Asn, Gly, Val, Pro or Lys); Xaa at res.56 = (Thr, Ala,
Val, Lys, Asp, Tyr, Ser, Gly, Ile or His); Xaa at res.57
= (Val, Ala or Ile); Xaa at res.58 = (Pro or Asp); Xaa
at res.59 = (Lys, Leu or Glu); Xaa at res.60 = (Pro, Val
20 or Ala); Xaa at res.63 = (Ala or Val); Xaa at res.65 =
(Thr, Ala or Glu); Xaa at res.66 = (Gln, Lys, Arg or
Glu); Xaa at res.67 = (Leu, Met or Val); Xaa at res.68 =
(Asn, Ser, Asp or Gly); Xaa at res.69 = (Ala, Pro or
Ser); Xaa at res.70 = (Ile, Thr, Val or Leu); Xaa at
25 res.71 = (Ser, Ala or Pro); Xaa at res.72 = (Val, Leu,
Met or Ile); Xaa at res.74 = (Tyr or Phe); Xaa at res.75
= (Phe, Tyr, Leu or His); Xaa at res.76 = (Asp, Asn or
Leu); Xaa at res.77 = (Asp, Glu, Asn, Arg or Ser); Xaa
at res.78 = (Ser, Gln, Asn, Tyr or Asp); Xaa at res.79 =
30 (Ser, Asn, Asp, Glu or Lys); Xaa at res.80 = (Asn, Thr
or Lys); Xaa at res.82 = (Ile, Val or Asn); Xaa at
res.84 = (Lys or Arg); Xaa at res.85 = (Lys, Asn, Gln,
His, Arg or Val); Xaa at res.86 = (Tyr, Glu or His); Xaa

at res.87 = (Arg, Gln, Glu or Pro); Xaa at res.88 =
 (Asn, Glu, Trp or Asp); Xaa at res.90 = (Val, Thr, Ala
 or Ile); Xaa at res.92 = (Arg, Lys, Val, Asp, Gln or
 Glu); Xaa at res.93 = (Ala, Gly, Glu or Ser); Xaa at
 5 res.95 = (Gly or Ala) and Xaa at res.97 = (His or Arg).

[0068] Generic Sequence 8 (SEQ ID NO: 6) includes all
 of Generic Sequence 7 and in addition includes the
 following sequence (SEQ ID NO: 9) at its N-terminus:

SEQ ID NO: 9

Cys	Xaa	Xaa	Xaa	Xaa
1				5

10 Accordingly, beginning with residue 7, each "Xaa" in
 Generic Sequence 8 is a specified amino acid defined as
 for Generic Sequence 7, with the distinction that each
 residue number described for Generic Sequence 7 is
 shifted by five in Generic Sequence 8. Thus, "Xaa at
 15 res.2 =(Tyr or Lys)" in Generic Sequence 7 refers to Xaa
 at res.7 in Generic Sequence 8. In Generic Sequence 8,
 Xaa at res.2 = (Lys, Arg, Ala or Gln); Xaa at res.3 =
 (Lys, Arg or Met); Xaa at res.4 = (His, Arg or Gln); and
 Xaa at res.5 = (Glu, Ser, His, Gly, Arg, Pro, Thr, or
 20 Tyr).

[0069] In another embodiment, useful osteogenic
 proteins include those defined by Generic Sequences 9
 and 10, defined as follows.

[0070] Specifically, Generic Sequences 9 and 10 are
 25 composite amino acid sequences of the following
 proteins: human OP-1, human OP-2, human OP-3, human BMP-
 2, human BMP-3, human BMP-4, human BMP-5, human BMP-6,
 human BMP-8, human BMP-9, human BMP-10, human BMP-11,
 Drosophila 60A, Xenopus Vg-1, sea urchin UNIVIN, human
 30 CDMP-1 (mouse GDF-5), human CDMP-2 (mouse GDF-6, human

BMP-13), human CDMP-3 (mouse GDF-7, human BMP-12), mouse GDF-3, human GDF-1, mouse GDF-1, chicken DORSALIN, dpp, Drosophila SCREW, mouse NODAL, mouse GDF-8, human GDF-8, mouse GDF-9, mouse GDF-10, human GDF-11, mouse GDF-11, human BMP-15, and rat BMP3b. Like Generic Sequence 7, Generic Sequence 9 is a 97 amino acid sequence that accommodates the C-terminal six cysteine skeleton and, like Generic Sequence 8, Generic Sequence 10 is a 102 amino acid sequence which accommodates the seven cysteine skeleton.

Generic Sequence 9

Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
1				5					10
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Pro	Xaa	Xaa	Xaa
				15					20
Xaa	Xaa	Xaa	Xaa	Cys	Xaa	Gly	Xaa	Cys	Xaa
				25					30
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				35					40
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				45					50
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				55					60
Xaa	Cys	Xaa	Pro	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				65					70
Xaa	Xaa	Leu	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				75					80
Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa	Xaa
				85					90
Xaa	Xaa	Xaa	Cys	Xaa	Cys	Xaa			
				95					

wherein each Xaa is independently selected from a group of one or more specified amino acids defined as follows:

"res." means "residue" and Xaa at res.1 = (Phe, Leu or Glu); Xaa at res.2 = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu); Xaa at res.3 = (Val, Ile, Leu or Asp); Xaa at res.4 = (Ser, Asp, Glu, Asn or Phe); Xaa at res.5 = (Phe or Glu); Xaa at res.6 = (Arg, Gln, Lys, Ser, Glu,

Ala or Asn); Xaa at res.7 = (Asp, Glu, Leu, Ala or Gln);
Xaa at res.8 = (Leu, Val, Met, Ile or Phe); Xaa at res.9
= (Gly, His or Lys); Xaa at res.10 = (Trp or Met); Xaa
at res.11 = (Gln, Leu, His, Glu, Asn, Asp, Ser or Gly);
5 Xaa at res.12 = (Asp, Asn, Ser, Lys, Arg, Glu or His);
Xaa at res.13 = (Trp or Ser); Xaa at res.14 = (Ile or
Val); Xaa at res.15 = (Ile or Val); Xaa at res.16 =
(Ala, Ser, Tyr or Trp); Xaa at res. 18 = (Glu, Lys, Gln,
Met, Pro, Leu, Arg, His or Lys); Xaa at res.19 = (Gly,
10 Glu, Asp, Lys, Ser, Gln, Arg or Phe); Xaa at res.20 =
(Tyr or Phe); Xaa at res.21 = (Ala, Ser, Gly, Met, Gln,
His, Glu, Asp, Leu, Asn, Lys or Thr); Xaa at res.22 =
(Ala or Pro); Xaa at res.23 = (Tyr, Phe, Asn, Ala or
Arg); Xaa at res.24 = (Tyr, His, Glu, Phe or Arg); Xaa
15 at res.26 = (Glu, Asp, Ala, Ser, Tyr, His, Lys, Arg, Gln
or Gly); Xaa at res.28 = (Glu, Asp, Leu, Val, Lys, Gly,
Thr, Ala or Gln); Xaa at res.30 = (Ala, Ser, Ile, Asn,
Pro, Glu, Asp, Phe, Gln or Leu); Xaa at res.31= (Phe,
Tyr, Leu, Asn, Gly or Arg); Xaa at res.32 = (Pro, Ser,
20 Ala or Val); Xaa at res.33 = (Leu, Met, Glu, Phe or
Val); Xaa at res.34 = (Asn, Asp, Thr, Gly, Ala, Arg, Leu
or Pro); Xaa at res.35 = (Ser, Ala, Glu, Asp, Thr, Leu,
Lys, Gln or His); Xaa at res.36 = (Tyr, His, Cys, Ile,
Arg, Asp, Asn, Lys, Ser, Glu or Gly); Xaa at res.37 =
25 (Met, Leu, Phe, Val, Gly or Tyr); Xaa at res.38 = (Asn,
Glu, Thr, Pro, Lys, His, Gly, Met, Val or Arg); Xaa at
res.39 = (Ala, Ser, Gly, Pro or Phe); Xaa at res.40 =
(Thr, Ser, Leu, Pro, His or Met); Xaa at res.41 = (Asn,
Lys, Val, Thr or Gln); Xaa at res.42 = (His, Tyr or
30 Lys); Xaa at res.43 = (Ala, Thr, Leu or Tyr); Xaa at
res.44 = (Ile, Thr, Val, Phe, Tyr, Met or Pro); Xaa at
res.45 = (Val, Leu, Met, Ile or His); Xaa at res.46 =
(Gln, Arg or Thr); Xaa at res.47 = (Thr, Ser, Ala, Asn

or His); Xaa at res.48 = (Leu, Asn or Ile); Xaa at
res.49 = (Val, Met, Leu, Pro or Ile); Xaa at res.50 =
(His, Asn, Arg, Lys, Tyr or Gln); Xaa at res.51 = (Phe,
Leu, Ser, Asn, Met, Ala, Arg, Glu, Gly or Gln); Xaa at
5 res.52 = (Ile, Met, Leu, Val, Lys, Gln, Ala or Tyr); Xaa
at res.53 = (Asn, Phe, Lys, Glu, Asp, Ala, Gln, Gly, Leu
or Val); Xaa at res.54 = (Pro, Asn, Ser, Val or Asp);
Xaa at res.55 = (Glu, Asp, Asn, Lys, Arg, Ser, Gly, Thr,
Gln, Pro or His); Xaa at res.56 = (Thr, His, Tyr, Ala,
10 Ile, Lys, Asp, Ser, Gly or Arg); Xaa at res.57 = (Val,
Ile, Thr, Ala, Leu or Ser); Xaa at res.58 = (Pro, Gly,
Ser, Asp or Ala); Xaa at res.59 = (Lys, Leu, Pro, Ala,
Ser, Glu, Arg or Gly); Xaa at res.60 = (Pro, Ala, Val,
Thr or Ser); Xaa at res.61 = (Cys, Val or Ser); Xaa at
15 res.63 = (Ala, Val or Thr); Xaa at res.65 = (Thr, Ala,
Glu, Val, Gly, Asp or Tyr); Xaa at res.66 = (Gln, Lys,
Glu, Arg or Val); Xaa at res.67 = (Leu, Met, Thr or
Tyr); Xaa at res.68 = (Asn, Ser, Gly, Thr, Asp, Glu, Lys
or Val); Xaa at res.69 = (Ala, Pro, Gly or Ser); Xaa at
20 res.70 = (Ile, Thr, Leu or Val); Xaa at res.71 = (Ser,
Pro, Ala, Thr, Asn or Gly); Xaa at res.72 = (Val, Ile,
Leu or Met); Xaa at res.74 = (Tyr, Phe, Arg, Thr, Tyr or
Met); Xaa at res.75 = (Phe, Tyr, His, Leu, Ile, Lys, Gln
or Val); Xaa at res.76 = (Asp, Leu, Asn or Glu); Xaa at
25 res.77 = (Asp, Ser, Arg, Asn, Glu, Ala, Lys, Gly or
Pro); Xaa at res.78 = (Ser, Asn, Asp, Tyr, Ala, Gly,
Gln, Met, Glu, Asn or Lys); Xaa at res.79 = (Ser, Asn,
Glu, Asp, Val, Lys, Gly, Gln or Arg); Xaa at res.80 =
(Asn, Lys, Thr, Pro, Val, Ile, Arg, Ser or Gln); Xaa at
30 res.81 = (Val, Ile, Thr or Ala); Xaa at res.82 = (Ile,
Asn, Val, Leu, Tyr, Asp or Ala); Xaa at res.83 = (Leu,
Tyr, Lys or Ile); Xaa at res.84 = (Lys, Arg, Asn, Tyr,
Phe, Thr, Glu or Gly); Xaa at res.85 = (Lys, Arg, His,

Gln, Asn, Glu or Val); Xaa at res.86 = (Tyr, His, Glu or Ile); Xaa at res.87 = (Arg, Glu, Gln, Pro or Lys); Xaa at res.88 = (Asn, Asp, Ala, Glu, Gly or Lys); Xaa at res.89 = (Met or Ala); Xaa at res.90 = (Val, Ile, Ala, Thr, Ser or Lys); Xaa at res.91 = (Val or Ala); Xaa at res.92 = (Arg, Lys, Gln, Asp, Glu, Val, Ala, Ser or Thr); Xaa at res.93 = (Ala, Ser, Glu, Gly, Arg or Thr); Xaa at res.95 = (Gly, Ala or Thr); Xaa at res.97 = (His, Arg, Gly, Leu or Ser). Further, after res.53 in rBMP3b and mGDF-10 there is an Ile; after res.54 in GDF-1 there is a T; after res.54 in BMP3 there is a V; after res.78 in BMP-8 and Dorsalin there is a G; after res.37 in hGDF-1 there is Pro, Gly, Gly, Pro.

[0071] Generic Sequence 10 (SEQ ID NO: 8) includes all of Generic Sequence 9 (SEQ ID NO: 7) and in addition includes the following sequence (SEQ ID NO: 9) at its N-terminus:

SEQ ID NO: 9

Cys	Xaa	Xaa	Xaa	Xaa
1				5

Accordingly, beginning with residue 6, each "Xaa" in Generic Sequence 10 is a specified amino acid defined as for Generic Sequence 9, with the distinction that each residue number described for Generic Sequence 9 is shifted by five in Generic Sequence 10. Thus, "Xaa at res.1 = (Tyr, Phe, His, Arg, Thr, Lys, Gln, Val or Glu)" in Generic Sequence 9 refers to Xaa at res.6 in Generic Sequence 10. In Generic Sequence 10, Xaa at res.2 = (Lys, Arg, Gln, Ser, His, Glu, Ala, or Cys); Xaa at res.3 = (Lys, Arg, Met, Lys, Thr, Leu, Tyr, or Ala); Xaa

at res.4 = (His, Gln, Arg, Lys, Thr, Leu, Val, Pro, or Tyr); and Xaa at res.5 = (Gln, Thr, His, Arg, Pro, Ser, Ala, Gln, Asn, Tyr, Lys, Asp, or Leu).

[0072] As noted above, certain currently preferred bone morphogenic polypeptide sequences useful in this invention have greater than 60% identity, preferably greater than 65% identity, with the amino acid sequence defining the preferred reference sequence of hOP-1. These particularly preferred sequences include allelic and phylogenetic counterpart variants of the OP-1 and OP-2 proteins, including the Drosophila 60A protein. Accordingly, in certain particularly preferred embodiments, useful BMPs include active proteins comprising pairs of polypeptide chains within the generic amino acid sequence herein referred to as "OPX" (SEQ ID NO: 4), which defines the seven cysteine skeleton and accommodates the homologies between several identified variants of OP-1 and OP-2. As described therein, each Xaa at a given position independently is selected from the residues occurring at the corresponding position in the C-terminal sequence of mouse or human OP-1 or OP-2.

SEQ ID NO: 4

25	Cys	Xaa	Xaa	His	Glu	Leu	Tyr	Val	Ser	Phe	Xaa	Asp	Leu	Gly	Trp	Xaa	Asp	Trp	1	5	10	15	
	Xaa	Ile	Ala	Pro	Xaa	Gly	Tyr	Xaa	Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Xaa	Phe	Pro	20	25	30	35
	Leu	Xaa	Ser	Xaa	Met	Asn	Ala	Thr	Asn	His	Ala	Ile	Xaa	Gln	Xaa	Leu	Val	His	Xaa	40	45	50	55
30	Xaa	Xaa	Pro	Xaa	Xaa	Val	Pro	Lys	Xaa	Cys	Cys	Ala	Pro	Thr	Xaa	Leu	Xaa	Ala		60	65	70	
	Xaa	Ser	Val	Leu	Tyr	Xaa	Asp	Xaa	Ser	Xaa	Asn	Val	Ile	Leu	Xaa	Lys	Xaa	Arg		75	80	85	90
35	Asn	Met	Val	Val	Xaa	Ala	Cys	Gly	Cys	His										95	100		

wherein Xaa at res.2 = (Lys or Arg); Xaa at res.3 = (Lys or Arg); Xaa at res.11 = (Arg or Gln); Xaa at res.16 = (Gln or Leu); Xaa at res.19 = (Ile or Val); Xaa at
5 res.23 = (Glu or Gln); Xaa at res.26 = (Ala or Ser); Xaa at res.35 = (Ala or Ser); Xaa at res.39 = (Asn or Asp); Xaa at res.41 = (Tyr or Cys); Xaa at res.50 = (Val or Leu); Xaa at res.52 = (Ser or Thr); Xaa at res.56 = (Phe or Leu); Xaa at res.57 = (Ile or Met); Xaa at res.58 =
10 (Asn or Lys); Xaa at res.60 = (Glu, Asp or Asn); Xaa at res.61 = (Thr, Ala or Val); Xaa at res.65 = (Pro or Ala); Xaa at res.71 = (Gln or Lys); Xaa at res.73 = (Asn or Ser); Xaa at res.75 = (Ile or Thr); Xaa at res.80 = (Phe or Tyr); Xaa at res.82 = (Asp or Ser); Xaa at
15 res.84 = (Ser or Asn); Xaa at res.89 = (Lys or Arg); Xaa at res.91 = (Tyr or His); and Xaa at res.97 = (Arg or Lys).

[0073] In still another preferred embodiment, useful BMPs have polypeptide chains with amino acid sequences
20 comprising a sequence encoded by a nucleic acid that hybridizes, under low, medium or high stringency hybridization conditions, to DNA or RNA encoding reference BMP sequences, e.g., C-terminal sequences defining the conserved seven cysteine domains of OP-1,
25 OP-2, BMP-2, BMP-4, BMP-5, BMP-6, 60A, GDF-5, GDF-6, GDF-7 and the like. As used herein, high stringent hybridization conditions are defined as hybridization according to known techniques in 40% formamide, 5 X SSPE, 5 X Denhardt's Solution, and 0.1% SDS at 37°C
30 overnight, and washing in 0.1 X SSPE, 0.1% SDS at 50°C. Standard stringent conditions are well characterized in commercially available, standard molecular cloning texts. See, for example, *Molecular Cloning A Laboratory*

Manual, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1989); *DNA Cloning*, Volumes I and II (D.N. Glover ed., 1985); *Oligonucleotide Synthesis* (M.J. Gait ed., 1984); *Nucleic Acid Hybridization* (B. D. Hames & S.J. Higgins eds. 1984); and B. Perbal, *A Practical Guide To Molecular Cloning* (1984), the disclosures of which are incorporated herein by reference.

[0074] As noted above, proteins useful in the present invention generally are dimeric proteins comprising a folded pair of the above polypeptides. In some embodiments, the pair of polypeptides are not disulfide bonded. In some embodiments the pair of polypeptides are disulfide bonded. Such disulfide bonded BMPs are inactive when reduced, but are active as oxidized homodimers and when oxidized in combination with others of this invention to produce heterodimers. Thus, members of a folded pair of bone morphogenic polypeptides in a morphogenically active protein can be selected independently from any of the specific polypeptides mentioned above.

[0075] The BMPs useful in the materials and methods of this invention include proteins comprising any of the polypeptide chains described above, whether isolated from naturally-occurring sources, or produced by recombinant DNA or other synthetic techniques, and includes allelic and phylogenetic counterpart variants of these proteins, as well as muteins thereof, and various truncated and fusion constructs. Deletion or addition mutants also are envisioned to be active, including those which may alter the conserved C-terminal six or seven cysteine domain, provided that the alteration does not functionally disrupt the

relationship of these cysteines in the folded structure. Accordingly, such active forms are considered the equivalent of the specifically described constructs disclosed herein. The proteins may include forms having
5 varying glycosylation patterns, varying N-termini, a family of related proteins having regions of amino acid sequence homology, and active truncated or mutated forms of native or biosynthetic proteins, produced by expression of recombinant DNA in host cells.

10 [0076] The BMPs contemplated herein can be expressed from intact or truncated cDNA or from synthetic DNAs in prokaryotic or eukaryotic host cells, and purified, cleaved, refolded, and dimerized to form morphogenically active compositions. Alternatively, cells expressing
15 recombinant may be used in the methods of this invention. Currently preferred host cells include, without limitation, prokaryotes including *E. coli* or eukaryotes including yeast, or mammalian cells, such as CHO, COS or BSC cells. One of ordinary skill in the art
20 will appreciate that other host cells can be used to advantage. Detailed descriptions of the bone morphogenic proteins useful in the practice of this invention, including how to make, use and test them for osteogenic activity, are disclosed in numerous
25 publications, including U.S. Patent Nos. 5,266,683 and 5,011,691, the disclosures of which are incorporated by reference herein.

[0077] Thus, in view of this disclosure and the knowledge available in the art, skilled genetic
30 engineers can isolate genes from cDNA or genomic libraries of various different biological species, which encode appropriate amino acid sequences, or construct DNAs from oligonucleotides, and then can express them in

various types of host cells, including both prokaryotes and eukaryotes, to produce large quantities of active proteins.

Serine/Threonine Kinase Receptors

5 **[0078]** TGF- β superfamily members elicit their cellular responses through formation of heteromeric complexes of specific type I and type II serine/threonine kinase receptors. To date five type II receptors and seven type I receptors (also termed
10 activin receptor-like kinases (ALKs) have been identified (See e.g., Derynck et al., Biochem. Biophys. Acta, 1333, pp. F105-F150 (1997); Massague, Annu. Rev. Biochem., 67, pp. 753-791 (1998)). The type II receptors are kinases and are constitutively active.
15 Upon ligand-mediated heteromeric complex formation, the type II receptor phosphorylates particular serine and threonine residues in the type I receptor juxtamembrane region, thereby activating the type I receptor.
 [0079] ALK-1 was identified as an endothelial
20 specific TGF- β type I receptor (see e.g., Oh et al., Proc. Natl. Acad. Sci. U.S.A., 97, pp. 2626-2631 (2000); see also, Lux et al., J. Biol. Chem., 274, 9984-9992 (1999); US patent 5,968,752). ALK-2 was identified as a type I receptor for activin, TGF- β and BMPs (see, e.g.,
25 Attisano et al., Cell, 75, pp. 671-680 (1993); Miettinen et al., J. Cell Biol., 127, pp. 2021-2036 (1994); ten Dijke et al., Science 264, 101-104 (1994); Macias-Silva et al., J. Cell Biol., 273, pp. 25628-25636 (1998); US Patent 6,271,365). ALK-4 and ALK-5 were identified as
30 activin and TGF- β type I receptors and ALK-3 and ALK-6 were identified as BMP type I receptors (see, e.g., Oh et al., *supra* and Lux et al., *supra*; US Patents 6,271,365 and 6,207,814). ALK-7 was identified as the

type I receptor for Nodal (see, e.g., Reissman et al., Genes Dev., 15, pp. 2010-22 (2001); US patent 5,891,638).

[0080] In some embodiments of the invention, type II
5 receptors are used. In some embodiments of the invention, type I receptors are used. In yet other embodiments of the invention, both type I and type II receptors are used. In some embodiments, the type I receptor is selected from the group consisting of ALK-1,
10 ALK-2, ALK-3, ALK-4, ALK-5, ALK-6, ALK-7, amino acid sequence variants and homologs thereof, including species homologs thereof and fragments thereof. In a preferred embodiment, the type I receptor is ALK-2, ALK-3 or ALK-6. In some embodiments, the type I and type II
15 receptors are recombinant.

[0081] Publications disclosing these sequences as well as their chemical and physical properties include: ALK-1 (US patent 5,968,752); ALK-2, ALK-4 and ALK-5 (US Patent 6,271,365); ALK-3 and ALK-6 (US Patent
20 6,207,814); ALK-7 (US patents 5,891,638, 5,614,609 and 5,789,565). These publications are incorporated herein by reference.

[0082] The type I and type II receptors contemplated herein can be expressed from intact or truncated cDNA or
25 from synthetic DNAs in prokaryotic or eukaryotic host cells (see discussion of protein expression, *infra*).

[0083] Thus, in view of this disclosure and the knowledge available in the art, skilled genetic engineers can express these receptors in various types
30 of host cells, including both prokaryotes and eukaryotes.

Smads

[0084] The Smad family of proteins can be divided into three subfamilies: R-Smads, Co-Smads and I-Smads. The subfamily of R-Smads can be divided into two groups: BMP-Smads and TGF- β /activin-Smads. Smad1, Smad5, and Smad8 are phosphorylated by ALK-1, ALK-2, ALK-3 and ALK-6, and Smad2 and Smad3 are activated by ALK-5 and ALK-4, respectively. Smad2 and Smad3 are also activated by ALK-7 (see e.g., Derynck et al., Biochim. Biophys. Acta, 1333, pp. F105-F150 (1997), Massague, Annu. Rev. Biochem., 67, 753-792 (1998); Itoh et al., Eur. J. Biochem., 267, 6954-6967 (2000)). The phosphorylated R-Smads form heteromeric complexes with Co-Smads. To date only one Co-Smad has been identified: Smad4. I-Smads, i.e., Smad6 and Smad7 inhibit TGF- β family signaling by preventing the activation of R-Smads and Co-Smads (see, e.g., Itoh et al., *supra*).

[0085] In some embodiments, the Smads used in the methods of this invention are selected from the group consisting of Smad1, Smad2, Smad3, Smad5, Smad8, amino acid sequence variants and homologs thereof, including species homologs thereof and fragments thereof. In a preferred embodiment, the Smad is Smad5. In another embodiment, the Smad is recombinant.

[0086] The sequences for Smads are included in the following publications or are readily available to the skilled worker from GenBank: Smad1 (rat: GenBank Accession No. AF067727; SEQ ID NO: 12 and 13); Smad2 (mouse: GenBank Accession No. NM_010754; SEQ ID NO: 14 and 15), Smad3 (human: GenBank Accession No. NM_005902; SEQ ID NO: 16 and 17), Smad4 (mouse: Genbank Accession No. NM_008540; human: GenBank Accession No. NM_005359; SEQ ID NO: 18 and 19), Smad5 (rat: GenBank Accession No.

NM_021692; SEQ ID NO: 20 and 21), Smad6 (US patents 6,534,476 and 6,270,994; human: GenBank Accession No. NM_005585; SEQ ID NO: 22 and 23), Smad7 (US patents 6,020,464 and 6,251,628; human: GenBank Accession No. NM_005904; SEQ ID NO: 24 and 25), and Smad8 (rat: GenBank Accession No. AF012347; SEQ ID NO: 26 and 27). These publications and GenBank Accession numbers are incorporated herein by reference.

[0087] The Smads contemplated herein can be expressed from intact or truncated cDNA or from synthetic DNAs in prokaryotic or eukaryotic host cells (see discussion of protein expression, *infra*).

[0088] Thus, in view of this disclosure and the knowledge available in the art, skilled genetic engineers can express these receptors in various types of host cells, including both prokaryotes and eukaryotes (e.g., progenitor cells).

Production or expression of BMPs, serine/threonine kinase receptors and Smads

[0089] The BMPs, serine/threonine kinase receptors and Smads according to this invention may be derived from a variety of sources. They may be isolated from natural sources, or may be produced by expressing an appropriate recombinant DNA molecule in a host cell. In addition, the BMPs, serine/threonine kinase receptors and Smads of this invention may be derived synthetically and synthetic proteins may optionally be expressed from a recombinant DNA molecule in a host cell.

1. Naturally-derived proteins

[0090] In one embodiment of this invention, the BMPs, serine/threonine kinase receptors and Smads are isolated from natural sources. BMPs, serine/threonine kinase

receptors and Smads may be purified from tissue sources, preferably mammalian tissue sources, using conventional physical and chemical separation techniques well known to those of skill in the art.

5 2. Recombinantly-expressed proteins

[0091] In another embodiment of this invention, the Smads, BMPs and serine/threonine kinase receptors and are produced by the expression of an appropriate recombinant DNA molecule in a host cell. The DNA and
10 amino acid sequences of Smads, BMPs and serine/threonine kinase receptors have been reported, and methods for their recombinant production are published and otherwise known to those of skill in the art. For a general discussion of cloning and recombinant DNA technology,
15 see Ausubel et al., *supra*; see also Watson et al., *Recombinant DNA*, 2d ed. 1992 (W.H. Freeman and Co., New York).

[0092] For cloning and expressing Smads, BMPs and serine/threonine kinase receptors, standard recombinant
20 DNA techniques may be used. With the DNA sequence available, a DNA fragment encoding any of these proteins be inserted into an expression vector selected to work in conjunction with a desired host expression system. The DNA fragment is cloned into the vector with the
25 proper transcription control elements. In some embodiments, the expression of the desired protein may be constitutive. In some embodiments, the expression of the desired protein is under the control of an inducible promoter.

30 Vectors

[0093] In some embodiments, the invention provides vectors comprising the nucleic acids encoding Smads, serine/threonine kinase receptors and/or BMPs. The

choice of vector and expression control sequences to which the nucleic acids of this invention are operably linked depends on the functional properties desired, e.g., protein expression, and the host cell to be transformed.

[0094] Expression control elements useful for regulating the expression of an operably linked coding sequence are known in the art. Examples include, but are not limited to, inducible promoters, constitutive promoters, secretion signals, and other regulatory elements. When an inducible promoter is used, it can be controlled, e.g., by a change in nutrient status (e.g. concentration of growth factors or BMPs), or a change in temperature, in the host cell medium.

[0095] An appropriate vector is selected according to the host system selected. Useful vectors include but are not limited to plasmids, cosmids, bacteriophage, insect and animal viral vectors, including retroviruses, and other single and double-stranded DNA viruses.

[0096] In some embodiments, it may be preferable to recombinantly produce a mammalian protein for therapeutic uses in mammalian cell culture systems in order to produce a protein whose structure resembles more closely that of the natural material. Recombinant protein production in mammalian cells requires the establishment of appropriate cells and cell lines that are easy to transfect, are capable of stably maintaining foreign DNA with an unarranged sequence, and which have the necessary cellular components for efficient transcription, translation, post-translational modification and secretion of the protein. In addition, a suitable vector carrying the gene of interest is necessary.

[0097] DNA vector design for transfection into mammalian cells should include appropriate sequences to promote expression of the gene of interest, including: appropriate transcription initiation, termination and enhancer sequences; efficient RNA processing signals such as splicing and polyadenylation signals; sequences that stabilize cytoplasmic mRNA; sequences that enhance translation efficiency (i.e., Kozak consensus sequence); sequences that enhance protein stability; and when desired, sequences that enhance protein secretion.

[0098] Preferred DNA vectors also include a marker gene and means for amplifying the copy number of the gene of interest. DNA vectors may also comprise stabilizing sequences (e.g., ori- or ARS-like sequences and telomere-like sequences), or may alternatively be designed to favor directed or non-directed integration into the host cell genome.

[0099] Substantial progress in the development of mammalian cell expression systems has been made in the last decade and many aspects of the system are well characterized. A detailed review of the production of foreign proteins in mammalian cells, including useful cells, protein expression-promoting sequences, marker genes, and gene amplification methods, is disclosed in M. M. Bendig, *Genetic Engineering*, 7, pp. 91-127 (1988).

[0100] Particular details of the transfection, expression and purification of recombinant proteins are well documented and are understood by those of skill in the art. Further details on the various technical aspects of each of the steps used in recombinant production of foreign genes in mammalian cell expression systems can be found in a number of texts and laboratory manuals in the art. See, e.g., F. M. Ausubel et al.,

ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York (1989).

[0101] Briefly, among the best characterized transcription promoters useful for expressing a foreign
5 gene in a particular mammalian cell are the SV40 early promoter, the adenovirus major late promoter (AdMLP), the mouse metallothionein-I promoter (mMT-I), the Rous sarcoma virus (RSV) long terminal repeat (LTR), the mouse mammary tumor virus long terminal repeat
10 (MMTV-LTR), and the human cytomegalovirus major intermediate-early promoter (hCMV). The DNA sequences for all of these promoters are known in the art and are available commercially.

[0102] One method of gene amplification in mammalian
15 cell systems is the use of the selectable dihydrofolate reductase (DHFR) gene in a dhfr- cell line. Generally, the DHFR gene is provided on the vector carrying the gene of interest, and addition of increasing concentrations of the cytotoxic drug methotrexate (MTX)
20 leads to amplification of the DHFR gene copy number, as well as that of the physically-associated gene of interest. DHFR as a selectable, amplifiable marker gene in transfected chinese hamster ovary cell lines (CHO cells) is particularly well characterized in the art.
25 Other useful amplifiable marker genes include the adenosine deaminase (ADA) and glutamine synthetase (GS) genes.

[0103] In one expression system, gene amplification is further enhanced by modifying marker gene expression
30 regulatory sequences (e.g., enhancer, promoter, and transcription or translation initiation sequences) to reduce the levels of marker protein produced. Lowering the level of DHFR transcription increases the DHFR gene

copy number (and the physically-associated gene) to enable the transfected cell to adapt to growth in even low levels of methotrexate (e.g., 0.1 μ M MTX). As will be appreciated by those skilled in the art, other useful
5 weak promoters, different from those disclosed and preferred herein, can be constructed using standard vector construction methodologies. In addition, other, different regulatory sequences also can be modified to achieve the same effect.

10 [0104] Another gene amplification scheme relies on the temperature sensitivity (ts) of BSC40-tsA58 cells transfected with an SV40 vector. Temperature reduction to 33 °C stabilizes the temperature sensitive SV40 T antigen, which leads to the excision and amplification
15 of the integrated transfected vector DNA thereby amplifying the physically associated gene of interest.

[0105] Eukaryotic cell expression vectors are known in the art and are commercially available. Typically, such vectors contain convenient restriction sites for
20 insertion of the desired DNA segment.

[0106] Eukaryotic cell expression vectors may include a selectable marker, e.g., a drug resistance gene. The neomycin phosphotransferase (neo) gene (Southern et al., 1982, J. Mol. Anal. Genet. 1:327-341) is an example of
25 such a gene.

[0107] To express the desired proteins of this invention, DNAs encoding the proteins (Smads, serine/threonine kinase receptors and/or BMPs are inserted into expression vectors such as plasmids,
30 retroviruses, cosmids, YACs, EBV-derived episomes, and the like. The expression vector and expression control sequences are chosen to be compatible with the expression host cell used. In some embodiments, Smad,

serine/threonine kinase receptors and BMP nucleic acids can be inserted into separate vectors. In some embodiments, the Smad, serine/threonine kinase receptor and BMP nucleic acids are inserted into the same
5 expression vector. In some embodiments the Smad and serine/threonine kinase receptor nucleic acids are inserted into the same vector. In some embodiments, the Smad and BMP nucleic acids are inserted into the same vector. Alternatively, any combination of Smad, BMP
10 and/or serine/threonine kinase receptor are inserted into the same vector.

[0108] A convenient vector is one that encodes a functionally complete protein. To the extent secretion of a desired protein is required, the recombinant
15 expression vector can also encode a signal peptide that facilitates secretion of the desired protein from a host cell.

[0109] Nucleic acid molecules encoding Smads, serine/threonine kinase receptors and/or BMPs, and
20 vectors comprising these nucleic acid molecules, can be used for transformation of a suitable host cell. Transformation can be by any suitable method. Methods for introduction of exogenous DNA into mammalian cells are well known in the art and include dextran-mediated
25 transfection, calcium phosphate precipitation, polybrene-mediated transfection, protoplast fusion, electroporation, encapsulation of the polynucleotide(s) in liposomes, and direct microinjection of the DNA into nuclei. In addition, nucleic acid molecules may be
30 introduced into mammalian cells by viral vectors.

[0110] Transformation of host cells can be accomplished by conventional methods suited to the vector and host cell employed. For transformation of

prokaryotic host cells, electroporation and salt treatment methods can be employed (Cohen et al., 1972, Proc. Natl. Acad. Sci. USA 69:2110-2114). For transformation of vertebrate cells, electroporation, cationic lipid or salt treatment methods can be employed. See, e.g., Graham et al., 1973, Virology 52:456-467; Wigler et al., 1979, Proc. Natl. Acad. Sci. USA 76:1373-1376f.

Host Cells

[0111] Host cells can be prokaryotic or eukaryotic. Useful host cells include but are not limited to bacteria such as *E. coli*, yeasts such as *Saccharomyces* and *Picia*, insect-baculovirus cell system, and primary, transformed or immortalized eukaryotic cells in culture. Preferred eukaryotic host cells include, but are not limited to, yeast and mammalian cells, e.g., Chinese hamster ovary (CHO) cell, NIH Swiss mouse embryo cells NIH-3T3, baby hamster kidney cells (BHK), C2C12 cells and BSC cells. Other useful eukaryotic cells include osteoprogenitor cells, cartilage progenitor cells, tendon progenitor cells, ligament progenitor cells and neural progenitor cells.

[0112] The methodology disclosed herein includes the use of COS cells for the rapid evaluation of vector construction and gene expression, and the use of established cell lines for long term protein production.

[0113] The choice of cells/cell lines is also important and depends on the needs of the skilled practitioner. Monkey kidney cells (COS) provide high levels of transient gene expression providing a useful means for rapidly testing vector construction and the expression of cloned genes. COS cells are transfected

with a simian virus 40 (SV40) vector carrying the gene of interest. The transfected COS cells eventually die, thus preventing the long term production of the desired protein product. However, transient expression does not
5 require the time consuming process required for the development of stable cell lines.

[0114] CHO cells are capable of successfully expressing a wide variety of proteins from a broad range of cell types. Thus, while the glycosylation pattern on
10 a recombinant protein produced in a mammalian cell expression system may not be identical to the natural protein, the differences in oligosaccharide side chains are often not essential for biological activity of the expressed protein.

15 [0115] The DHFR gene also may be used as part of a gene amplification scheme for CHO cells. Another gene amplification scheme relies on the temperature sensitivity (ts) of BSC40-tsA58 cells transfected with an SV40 vector. Temperature reduction to 33 °C
20 stabilizes the ts SV40 T antigen which leads to the excision and amplification of the integrated transfected vector DNA, thereby also amplifying the associated gene of interest.

[0116] Stable cell lines were established for CHO
25 cells as well as BSC40-tsA58 cells (hereinafter referred to as "BSC cells"). The various cells, cell lines and DNA sequences chosen for mammalian cell expression of the Smads, serine/threonine kinase receptors and BMPs of this invention are well characterized in the art and are
30 readily available. Other promoters, selectable markers, gene amplification methods and cells also may be used to express the Smads, serine/threonine kinase receptors and BMPs of this invention. Particular details of the

transfection, expression, and purification of recombinant proteins are well documented in the art and are understood by those having ordinary skill in the art. Further details on the various technical aspects of each of the steps used in recombinant production of foreign genes in mammalian cell expression systems can be found in a number of texts and laboratory manuals in the art. See, e.g., F. M. Ausubel et al., ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York (1989).

Progenitor cells

[0117] The progenitor cells that are induced to proliferate and/or differentiate in the present invention are preferably mammalian cells. Preferred progenitor cells include mammalian chondroblasts, osteoblasts and neuroblasts, all earlier developmental precursors thereof, and all cells that develop therefrom (e.g., chondroblasts, pre-chondroblasts and chondrocytes). However, any non-mammalian progenitor cells are also likely to be useful in the methods of the present invention. It is, thus, envisioned that when schemes become available for implanting xenogeneic cells into humans without causing adverse immunological reactions, non-mammalian progenitor cells will be useful for tissue regeneration and repair in humans.

[0118] In some embodiments, the progenitor cells comprise a nucleic acid encoding a Smad and optionally a nucleic acid encoding one or more serine/threonine kinase receptor and/or a nucleic acid encoding one or more BMP. In some embodiments, the progenitor cells comprise vectors comprising a nucleic acid encoding a Smad and optionally a nucleic acid encoding one or more

serine/threonine kinase receptor and/or a nucleic acid encoding one or more BMP. In some embodiments, the nucleic acid encoding Smad, serine/threonine kinase receptor and/or BMP are in one vector. In some
5 embodiments, the nucleic acid encoding Smad, serine/threonine kinase receptor and/or BMP are in different vectors. In some embodiments, the nucleic acids are recombinant.

[0119] In some embodiments, only type I
10 serine/threonine kinase receptors are used. In some embodiments, only type II serine/threonine kinase receptors are used. In other embodiments, both type I and type II serine/threonine kinase receptors are used. In some embodiments, more than one BMP will be
15 administered to the desired cell or tissue. In some embodiments, two BMPs will be used. In some embodiments, three BMPs will be used. The particular choice of combination of BMPs and the relative concentrations at which they are combined may be varied
20 systematically to optimize the tissue type induced at a selected treatment site using the procedures described herein. The relative concentrations of the BMPs either alone or in combination that will optimally induce tissue formation when administered to a mammal may be
25 determined empirically by the skilled practitioner using the procedures described herein.

Gene Therapy

[0120] The Smad, serine/threonine kinase receptor and/or BMP proteins can be produced *in vivo* in a mammal,
30 e.g., a human patient, using a gene therapy approach for inducing tissue formation, repairing a tissue defect or regenerating tissue at a target locus. This involves

administration of a suitable Smad, serine/threonine kinase receptor and/or BMP protein-encoding nucleic acid operably linked to suitable expression control sequences. Preferably, these sequences are incorporated
5 into a viral vector. Suitable viral vectors for such gene therapy include adenoviral vectors, lentiviral vectors, baculoviral vectors, Epstein Barr viral vectors, papovaviral vectors, vaccinia viral vectors, herpes simplex viral vectors, and adeno associated virus
10 (AAV) vectors. The viral vector can be a replication-defective viral vector. A preferred adenoviral vector has a deletion in its E1 gene or E3 gene. When an adenoviral vector is used, preferably the mammal is not exposed to a nucleic acid encoding a selectable marker
15 gene.

Formulations

[0121] The proteins (e.g., BMPs) of the present invention can be formulated as part of a pharmaceutical composition. The pharmaceutical compositions provided
20 by this invention comprise at least one protein. The compositions of this invention will be administered at an effective dose to induce the particular type of tissue at the treatment site selected according to the particular clinical condition addressed. Determination
25 of a preferred pharmaceutical formulation and a therapeutically effective dose regimen for a given application is well within the skill of the art taking into consideration, for example, the mode of administration, the condition and weight of the patient,
30 the extent of the desired treatment and the tolerance of the patient for the treatment.

[0122] Administration of the proteins of this invention, may be accomplished using any of the conventional modes of administration.

[0123] The pharmaceutical compositions comprising a protein of this invention may be in a variety of forms. These include, for example, solid, semi-solid and liquid dosage forms such as tablets, pills, powders, liquid solutions or suspensions, suppositories, and injectable and infusible solutions. The preferred form depends on the intended mode of administration and therapeutic application and may be selected by one skilled in the art. Modes of administration may include oral, parenteral, subcutaneous, intravenous, intralesional or topical administration. In most cases, the pharmaceutical compositions of this invention will be administered in the vicinity of the treatment site in need of tissue regeneration or repair.

[0124] The pharmaceutical compositions comprising a protein of this invention may, for example, be placed into sterile, isotonic formulations with or without cofactors which stimulate uptake or stability. The formulation is preferably liquid, or may be lyophilized powder. For example, the BMP of this invention may be diluted with a formulation buffer comprising 5.0 mg/ml citric acid monohydrate, 2.7 mg/ml trisodium citrate, 41 mg/ml mannitol, 1 mg/ml glycine and 1 mg/ml polysorbate 20. This solution can be lyophilized, stored under refrigeration and reconstituted prior to administration with sterile Water-For-Injection (USP).

[0125] The compositions also will preferably include conventional pharmaceutically acceptable carriers well known in the art (see for example Remington's Pharmaceutical Sciences, 16th Edition, 1980, Mac

Publishing Company). Such pharmaceutically acceptable carriers may include other medicinal agents, carriers, genetic carriers, adjuvants, excipients, etc., such as human serum albumin or plasma preparations. The

5 compositions are preferably in the form of a unit dose and will usually be administered as a dose regimen that depends on the particular tissue treatment.

[0126] The pharmaceutical compositions of this invention may also be administered in conjunction with a
10 morphogenic device using, for example, microspheres, liposomes, other microparticulate delivery systems or sustained release formulations placed in, near, or otherwise in communication with affected tissues or the bloodstream bathing those tissues.

15 [0127] Liposomes containing a protein of this invention can be prepared by well-known methods (See, e.g. DE 3,218,121; Epstein et al., Proc. Natl. Acad. Sci. U.S.A., 82, pp. 3688-92 (1985); Hwang et al., Proc. Natl. Acad. Sci. U.S.A., 77, pp. 4030-34 (1980); U.S.
20 Patent Nos. 4,485,045 and 4,544,545). Ordinarily the liposomes are of the small (about 200-800 Angstroms) unilamellar type in which the lipid content is greater than about 30 mol.% cholesterol. The proportion of cholesterol is selected to control the optimal rate of
25 BMP release.

[0128] The proteins of this invention may also be attached to liposomes containing other biologically active molecules such as immunosuppressive agents, cytokines, etc., to modulate the rate and
30 characteristics of tissue induction. Attachment of BMPs to liposomes may be accomplished by any known cross-linking agent such as heterobifunctional cross-linking agents that have been widely used to couple toxins or

chemotherapeutic agents to antibodies for targeted delivery. Conjugation to liposomes can also be accomplished using the carbohydrate-directed cross-linking reagent 4-(4-maleimidophenyl) butyric acid hydrazide (MPBH) (Duzgunes et al., J. Cell. Biochem. Abst. Suppl. 16E 77 (1992)).

[0129] The following are examples which illustrate the methods of this invention. These examples should not be construed as limiting: the examples are included for purposes of illustration and the present invention is limited only by the claims.

Example 1 - Effect of CDMP-1 and OP-1 on Smad5 levels

[0130] Total C2C12 cell lysates treated with solvent control (lane 1), CDMP-1 (200 ng/ml), or OP-1 (100 ng/ml) were collected after 5 days. Proteins in the lysates were resolved by SDS-PAGE, transferred to nylon membranes, and analyzed using the anti-human Smad5 antibody (Cell Signaling Technology, Inc., Beverly, MA), as the primary antibody and anti-rabbit IgG HRP-conjugated antibody was used as the secondary antibody. Immunoreactive bands were detected using the SuperSignal chemiluminescent detection system (Pierce, Rockford, IL), according to the manufacturer's instructions. Representative results of two independent experiments are shown. Western blot analyses showed that the Smad5 protein level in control C2C12 cells was hardly detectable by Western blot analysis (Figure 1, lane 1). CDMP-1 alone stimulated Smad5 protein expression in C2C12 cells (Figure 1, lane 2). OP-1 alone stimulated Smad5 protein expression to a higher level (Figure 1, lane 3). The combination of CDMP-1 and OP-1 significantly stimulated Smad5 protein expression even

more (Figure 1, lane 4). The current observation that the protein level of Smad5 was dramatically increased in cells treated with the combination of OP-1 and CDMP-1 would support the supposition that the effect of the combination of OP-1 and CDMP-1 is likely directed at the Smad signaling pathway.

Example 2 - Co-transfection with the OP-1 Gene and the CDMP-1 Gene

10 [0131] Target cells will be transfected with plasmids or viral carriers containing the OP-1 gene and the CDMP-1 gene under an appropriate promoter. In addition, these cells will be transfected with plasmids or viral carriers containing the Smad5 gene. For example, 15 plasmid pW24 (10.35 kb) that contains the coding sequence for OP-1 under the control of the CMV promoter may be used. Similarly, plasmids containing the CMDP-1 gene (pCDMP-1) or the Smad5 gene (pSmad5) will be constructed by replacing the OP-1 gene in the pW24 20 plasmid. Alternatively, these genes may be placed under the control of promoters that can be induced thus allowing control of expression at will. Additional vectors, such as adenoviral vectors may be constructed and tested in order to optimize and maximize expression.

25 [0132] Target cells that have been shown capable to become osteoblastic upon treatment with OP-1 include osteoblastic cells such as FRC cells, and pluripotent mesenchymal cells such as C2C12 and C3H10T1/2 cell lines. Cells of other orthopaedic utility may also be 30 used. These include cells derived from cartilage, tendon, ligament, and meniscus.

[0133] Transfection studies will be carried out using the calcium phosphate-DNA coprecipitation method or

FuGene6 (Roche Diagnostics). Briefly, cells will be transfected with the specified plasmid DNAs (1 µg/ml) in serum-free medium for 4-6 h. The medium will be replaced with complete αMEM containing 10% FBS.

5 **Example 3 - Effect of OP-1, CDMP-1 and/or Smad5 on Alkaline Phosphatase Activity**

[0134] The effect of co-transfection of FRC cells with pW24 and pCDMP-1 and/or pSmad5 will be examined. For example, FRC cells will be transfected with pW24, pCDMP-1 and/or pSmad5. After 48 h, total AP activity will be measured. It is anticipated that the AP activity in cells co-transfected with pW24, pCDMP-1 and/or pSmad5 will increase as a function of pCDMP-1 and/or pSmad5 concentration. The increase should be beyond that in cells transfected with pW24 alone. The AP activity in cells transfected with pCDMP-1 alone will also show a lesser but significant increase compared to the non-transfected control or cells transfected with the empty plasmids (vectors without the OP-1, CDMP-1 and/or Smad5 genes). Co-transfection with pW24 and the empty plasmid is not expected to result in an increase in AP activity beyond that by pW24 alone. Protein levels of OP-1, CDMP-1, and Smad5 under these conditions will be measured by Western blot analysis. Control experiments will include co-transfection of FRC cells with the empty plasmids, i.e. vector without the OP-1, CDMP-1 or Smad5 genes. It is expected that the AP activity will not be elevated beyond the non-transfected controls. The expression of additional osteoblastic cell markers, such as osteocalcin (OC) and bone sialoprotein (BSP) will also be monitored using either Northern blot analysis or real-time PCR. Transfection

conditions for other cell types will be optimized accordingly using art recognized methods.

Example 4 - Effect of OP-1, CDMP-1 and/or Smad5 on Mineralized Bone Nodule Formation

5 [0135] The effect of co-transfection of FRC cells with pW24 and pCDMP-1 or pSmad5 on mineralized bone nodule formation will be examined. Confluent FRC cells will be co-transfected with pW24 and pCDM-1 and/or pSmad5 using FuGene6 using the optimal ratio of pW24 to
10 pCDMP-1 or pSmad5 as determined by the experiments described above. Cells will be cultured in complete α MEM containing 5% FBS, ascorbic acid, and β -glycerol phosphate in the presence of 250 μ g/ml of Neomycin. Media will be changed every 3 days. Progress of nodule
15 formation will be monitored periodically and the images will be captured using an Olympus CK2 inverted microscope equipped with a CCD camera. At the termination of the experiments, cultures will be stained by Alizarin Red-S staining to assess the extent of
20 formation of mineralized bone nodules.

Example 5 - In vivo Expression of OP-1, CDMP-1 and Smad5

[0136] In vivo studies will be conducted using two experimental approaches: (i) Direct injection of OP-1
25 expressing vectors together with CDMP-1 and/or Smad5-expressing vectors into muscles of mice, and (ii) injection of transfected cells into muscles.

[0137] For direct injection experiments, nude mice will be injected with vectors expressing OP-1, CDMP-1
30 and/or Smad5 (pW24, pCDMP-1 and/or pSmad5, respectively) with a 27-gauge needle subcutaneously into a male homozygous nude mouse. Standard aseptic techniques will be used in all manipulations. To determine *in vivo*

osteogenic dose response of the vectors, eight mice will be used. Each mouse will be injected with 0.1 - 10 mg/ml vectors in 100 μ l each. Body weight and growth at the site of injection will be followed daily via in-life measurement of the mass. The cross-sectional area of the mass will be measured with a vernier caliper. The size of the mass will be calculated using the formula: length/2 x width/2 x π . The mass and the body weight will be plotted as a function of time following injection. The animals will be monitored for 49 days. At necropsy, the mass at the site of injection will be collected, fixed, stained with hematoxylin and eosin, and subjected to histological analysis. Controls will include mice injected with individual pW24, pCDMP-1 and/or pSmad5 alone. It is anticipated that the bone mass in mice injected with the combination of the pW24 and pCDMP-1, or pW24 plus pSmad5 will be greater than that injected with individual vector alone.

[0138] For experiments using injection of cells, similar experiments as described above will be conducted except that animals will be injected with cells co-transfected with vectors carrying the OP-1, CDMP-1, and/or Smad5 genes. Accordingly, cells will be grown to mid-log phase and transfected with a combination of vectors expressing OP-1, CDMP-1 and/or Smad5 as described above using the optimal ratio of the two vectors. Cells will be removed from the culturing dishes by trypsin-EDTA digestion. Trypsin will be inactivated by serum (10%) and removed by repeated washings with HBSS. Cells will be suspended in a minimal volume of HBSS and injected with a 27-gauge needle subcutaneously into the flank of a male homozygous nude mouse. Standard aseptic techniques will

be used in all manipulations. Eight nude mice will be injected with 10^6 cells in 100 μ l each. Outcome measurements as described above will be conducted. It is anticipated that the bone mass in mice injected with cells transfected with the combination of the pW24 and pCDMP-1, or pW24 plus pSmad5 will be greater than that injected with cells transfected with individual vector alone.

Example 6 - Gene Therapy In Patients Using Transfected

Cells

[0139] For cell therapeutics with tranfected genes, appropriate cells will be transfected *in vitro* with DNA vectors carrying the OP-1 gene, the CDMP-1 gene, and/or the Smad5 gene and optionally a serine/threonine kinase receptor. Appropriate cells include osteoblasts or osteoblastic cell progenitors for the repair of bone defects. For repair of cartilage regeneration, cells of chondrocyte origin or chondrogenitor cells will be appropriate. Similarly, for the regeneration of tendons or ligaments, the appropriate cells include progenitor cells of tendon or ligament origin. The transfected cells will be cultured to allow expression of the transfected gene(s). The cells will then be injected or implanted into a defect site in a patient. The defect site may be in bone, cartilage, tendon, ligament or neural tissue. The number of cells injected or implanted into the defect will depend on the size of the defect. Exemplary DNA vectors will be pW24, pCDMP-1, or pSmad5 as described previously.

Example 7 - Gene Therapy In Patients Using Transfected

Cells

[0140] For directed gene therapy, a combination of vectors as described above carrying the OP-1 gene, the

cDMP-1 gene, the Smad5 gene and/or a serine/threonine kinase will be injected into the defect site in a patient. The genes encoding each of the proteins may be placed in the same vector or in separate vectors.

5 **Example 8 - Monitoring Effects of Gene Therapy in Patients**

[0141] The repair site will be monitored radiographically every two weeks for a minimum of two years. It is anticipated that the defect site which
10 receives the combination of OP-1 + CDMP-1 + Smad or OP-1 + Smad5 (delivered by either of the methods described in Examples 6 and 7) will exhibit a faster rate of repair than that receives OP-1 alone.

[0142] In case of cartilage and tendon/ligament
15 repair, delivery of pCDMP-1 alone to a defect site will induce tissue formation. However, delivery of OP-1 + CDMP-1 to a defect site should exhibit a faster rate of repair than delivery of CDMP-1 or Smad5 alone. In all cases, the treated sites should exhibit regeneration
20 whereas those treated with vehicle should not.

We Claim:

1. A method of inducing the expression of a Smad in a cell or tissue comprising contacting the cell or tissue capable of expressing the Smad with a bone morphogenic protein.
2. The method according to claim 1, wherein the cell or tissue is contacted with two bone morphogenic proteins.
3. The method according to claim 1 or 2, wherein each bone morphogenic protein is independently selected from the group consisting of OP-1 (BMP-7), OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16,
5 BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A
10 protein, NODAL, UNIVIN, SCREW, ADMP, and NEURAL.
4. The method according to claim 1, wherein the bone morphogenic protein is OP-1.
5. The method according to claim 1, wherein the bone morphogenic protein is CDMP-1 or GDF-5.
6. The method according to claim 2, wherein the first bone morphogenic protein is OP-1 and the second bone morphogenic protein is selected from the group consisting of CDMP-1, and GDF-5.

7. The method according to claim 1 or 2, wherein the Smad is selected from the group consisting of Smad1, Smad2, Smad3, Smad5 and Smad8.

8. The method according to claim 7, wherein the Smad is Smad5.

9. The method according to claim 1, wherein the Smad is a recombinant Smad.

10. The method according to claim 1 or 2, wherein the cell or tissue is further capable of expressing a serine/threonine kinase receptor.

11. The method according to claim 10, wherein the serine/threonine kinase receptor is a type I receptor.

12. The method according to claim 10, wherein the serine/threonine kinase receptor is a type II receptor.

13. The method according to claim 10, wherein the cell or tissue is further capable of expressing both type I and type II serine/threonine kinase receptor.

14. The method according to claim 11 or 13, wherein the type I receptor is selected from the group consisting of ALK-1, ALK-2, ALK-3, ALK-4, ALK-5, ALK-6, and ALK-7.

15. The method according to claim 10 wherein the serine/threonine kinase receptor is a recombinant serine/threonine kinase receptor.

16. The method according to claim 1 or 2 wherein the tissue is selected from the group consisting of bone, cartilage, tendon, ligament and neural tissue.

17. The method according to claim 1 or 2 wherein the cell is a progenitor cell.

18. The method according to claim 17, wherein the progenitor cell is selected from the group consisting of an osteoprogenitor cell, a cartilage progenitor cell, a ligament progenitor cell, a tendon
5 progenitor cell, and a neural progenitor cell.

19. A method of inducing tissue formation at a target locus in a mammal comprising the step of administering to the target locus a nucleic acid encoding a Smad.

20. A method of inducing tissue formation at a target locus in a mammal comprising the step of administering to the target locus a vector comprising a nucleic acid encoding a Smad operably linked to an expression control sequence.

21. A method of inducing tissue formation at a target locus in a mammal comprising the step of administering to the target locus a cell comprising a vector comprising a nucleic acid encoding a Smad
5 operably linked to an expression control sequence.

22. A method of repairing a tissue defect or regenerating tissue at a target locus in a mammal comprising the step of administering to the target locus a nucleic acid encoding a Smad.

23. A method of repairing a tissue defect or regenerating tissue at a target locus in a mammal comprising the step of administering to the target locus a vector comprising a nucleic acid encoding a Smad operably linked to an expression control sequence.

24. A method of repairing a tissue defect or regenerating tissue at a target locus in a mammal comprising the step of administering to the target locus a cell comprising a vector comprising a nucleic acid encoding a Smad operably linked to an expression control sequence.

25. The method according to any one of claims 20, 21, 23 or 24, wherein the expression control sequence comprises a constitutive promoter.

26. The method according to claim 20, 21, 23 or 24, wherein the expression control sequence comprises an inducible promoter.

27. The method according to any one of claims 19-24, wherein the Smad is selected from the group consisting of Smad1, Smad2, Smad3, Smad5 and Smad8.

28. The method according to claim 27, wherein the Smad is Smad5.

29. The method according to any one of claims 19-28, wherein the Smad is recombinant Smad.

30. The method according to any one of claims 19-24 further comprising the step of

administering to the target locus a nucleic acid encoding a serine/threonine kinase receptor.

31. The method according to any one of claims 19-24 further comprising the step of administering to the target locus a vector comprising a nucleic acid encoding a serine/threonine kinase receptor operably linked to an expression control sequence.

32. The method according to any one of claims 19-24 further comprising the step of administering to the target locus a cell comprising a vector comprising a nucleic acid encoding a serine/threonine kinase receptor operably linked to an expression control sequence.

33. The method according to claim 31 or 32, wherein the expression control sequence operably linked to the serine/threonine kinase receptor comprises a constitutive promoter.

34. The method according to claim 31 or 32, wherein the expression control sequence operably linked to the serine/threonine kinase receptor comprises an inducible promoter.

35. The method according to any one of claims 33-32, wherein the serine/threonine kinase receptor is a type I receptor.

36. The method according to any one of claims 33-32, wherein the serine/threonine kinase receptor is a type II receptor.

37. The method according to any one of claims 33-32, wherein both type I and type II serine/threonine kinase receptors are administered.

38. The method according to any one of claim 35 or 37, wherein the type I receptor is selected from the group consisting of ALK-1, ALK-2, ALK-3, ALK-4, ALK-5, ALK-6, and ALK-7.

39. The method according to claim 38, wherein the serine/threonine kinase receptor is selected from the group consisting of ALK-2, ALK-3, and ALK-6.

40. The method according to claim 30-32, wherein the serine/threonine kinase receptor is a recombinant ALK.

41. The method according to any one of claims 19-24, further comprising the step of administering to the target locus a bone morphogenic protein.

42. The method according to any one of claims 19-24, further comprising the step of administering to the target locus a nucleic acid encoding a bone morphogenic protein.

43. The method according to any one of claims 19-24, further comprising the step of administering to the target locus a vector comprising a nucleic acid encoding a bone morphogenic protein operably linked to an expression control sequence.

44. The method according to any one of claims 19-24, further comprising the step of administering to the target locus a cell comprising a vector comprising a nucleic acid encoding a bone morphogenic protein operably linked to an expression control sequence.

45. The method according to any one of claims 41-44, wherein the bone morphogenic protein is selected from the groups consisting of OP-1 (BMP-7), OP-2, OP-3, COP-1, COP-3, COP-4, COP-5, COP-7, COP-16, BMP-2, BMP-3, BMP-3b, BMP-4, BMP-5, BMP-6, BMP-9, BMP-10, BMP-11, CDMP-3 (BMP-12), CDMP-2 (BMP-13), CDMP-1 (BMP-14), BMP-15, BMP-16, BMP-17, BMP-18, GDF-1, GDF-2, GDF-3, GDF-5, GDF-6, GDF-7, GDF-8, GDF-9, GDF-10, GDF-11, GDF-12, MP121, dorsalin-1, DPP, Vg-1, Vgr-1, 60A protein, NODAL, UNIVIN, SCREW, ADMP, and NEURAL.


46. The method according to claim 45, wherein the bone morphogenic protein is OP-1.

47. The method according to any one of claims 19-24, wherein the tissue is selected from the group consisting of bone, cartilage, tendon, ligament and neural tissue.

48. The method according to claim 21 or 24, wherein the cell is a progenitor cell.

49. The method according to claim 48, wherein the progenitor cell is selected from the group consisting of an osteoprogenitor cell, a cartilage progenitor cell, a ligament progenitor cell, a tendon progenitor cell, and a neural progenitor cell.


1/2

FIGURE 1

2/2

Figure 2

Plasmid map of pW24 containing OP-1 gene



SEQUENCE LISTING

<110> Stryker Corporation

<120> THERAPEUTIC METHODS USING SMADs

<130> STK-15 PCT

<140> Not yet assigned

<141> 2005-02-04

<160> 27

<170> PatentIn version 3.2

<210> 1

<211> 431

<212> PRT

<213> Homo sapiens

<400> 1

Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
 1 5 10 15

Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
 20 25 30

Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
 35 40 45

Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
 50 55 60

Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
 65 70 75 80

Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly
 85 90 95

Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser
 100 105 110

Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr
 115 120 125

Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys
 130 135 140

Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu
 145 150 155 160

Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile
 165 170 175

Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Glu Arg Ile
 180 185 190

Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu
 195 200 205

Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu
 210 215 220

Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg
 225 230 235 240

His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser
 245 250 255

Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn
 260 265 270

Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe
 275 280 285

Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser
 290 295 300

Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu
 305 310 315 320

Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr
 325 330 335

Val Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu
 340 345 350

Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn
 355 360 365

Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His
 370 375 380

Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln
 385 390 395 400

Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile
 405 410 415

Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430

<210> 2
 <211> 96
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic amino acid sequence COP-5

<400> 2

Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asp Asp Trp Ile Val Ala
 1 5 10 15

Pro Pro Gly Tyr Gln Ala Phe Tyr Cys His Gly Glu Cys Pro Phe Pro
 20 25 30

Leu Ala Asp His Phe Asn Ser Thr Asn His Ala Val Val Gln Thr Leu
 35 40 45

Val Asn Ser Val Asn Ser Lys Ile Pro Lys Ala Cys Cys Val Pro Thr
 50 55 60

Glu Leu Ser Ala Ile Ser Met Leu Tyr Leu Asp Glu Asn Glu Lys Val
 65 70 75 80

Val Leu Lys Tyr Asn Gln Glu Met Val Val Glu Gly Cys Gly Cys Arg
 85 90 95

<210> 3
 <211> 96
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Synthetic amino acid sequence COP-7

<400> 3

Leu Tyr Val Asp Phe Ser Asp Val Gly Trp Asn Asp Trp Ile Val Ala
 1 5 10 15

Pro Pro Gly Tyr His Ala Phe Tyr Cys His Gly Glu Cys Pro Phe Pro
 20 25 30

Page 4

specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (23)..(23)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (26)..(26)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (35)..(35)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (39)..(39)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (41)..(41)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (50)..(50)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (52)..(52)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (56)..(58)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (60)..(61)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (65)..(65)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>

<221> misc_feature
<222> (71)..(71)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (73)..(73)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (75)..(75)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (80)..(80)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (82)..(82)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (84)..(84)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (87)..(87)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (89)..(89)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (91)..(91)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (97)..(97)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<400> 4

Cys Xaa Xaa His Glu Leu Tyr Val Xaa Phe Xaa Asp Leu Gly Trp Xaa
 1 5 10 15
 Asp Trp Xaa Ile Ala Pro Xaa Gly Tyr Xaa Ala Tyr Tyr Cys Glu Gly
 20 25 30
 Glu Cys Xaa Phe Pro Leu Xaa Ser Xaa Met Asn Ala Thr Asn His Ala
 35 40 45
 Ile Xaa Gln Xaa Leu Val His Xaa Xaa Xaa Pro Xaa Xaa Val Pro Lys
 50 55 60
 Xaa Cys Cys Ala Pro Thr Xaa Leu Xaa Ala Xaa Ser Val Leu Tyr Xaa
 65 70 75 80
 Asp Xaa Ser Xaa Asn Val Xaa Leu Xaa Lys Xaa Arg Asn Met Val Val
 85 90 95
 Xaa Ala Cys Gly Cys His
 100

<210> 5
 <211> 97
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Generic Sequence 7

<220>
 <221> MOD_RES
 <222> (1)..(97)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (2)..(4)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (6)..(8)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (11)..(16)
 <223> Xaa is independently selected from a group of one or more

specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (18)..(21)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (23)..(23)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (26)..(26)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (28)..(28)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (30)..(31)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (33)..(40)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (44)..(60)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (63)..(63)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (65)..(72)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (74)..(80)
<223> Xaa is independently selected from a group of one or more

specified amino acids as defined in the specification

```

<220>
<221> misc_feature
<222> (82)..(82)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (84)..(88)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (90)..(90)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (92)..(93)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (95)..(95)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (97)..(97)
<223> Xaa is independently selected from a group of one or more
specified amino acids as defined in the specification

<400> 5

Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa Xaa Xaa Xaa Xaa Xaa
1          5          10          15

Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly Xaa Cys Xaa Xaa Pro
          20          25          30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala Xaa Xaa Xaa Xaa Xaa
35          40          45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Cys Xaa Pro
50          55          60

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65          70          75          80

```

Xaa

<220>

<221> misc_feature
<222> (31)..(31)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (33)..(33)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (35)..(36)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (38)..(45)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (49)..(65)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (68)..(68)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (70)..(77)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (79)..(85)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (87)..(87)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (89)..(93)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>

<221> misc_feature
 <222> (95)..(95)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (97)..(98)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (100)..(100)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (102)..(102)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<400> 6

Cys Xaa Xaa Xaa Xaa Leu Xaa Xaa Xaa Phe Xaa Xaa Xaa Gly Trp Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Ala Xaa Tyr Cys Xaa Gly
 20 25 30

Xaa Cys Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Asn His Ala
 35 40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 50 55 60

Xaa Cys Cys Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Leu Xaa Xaa
 65 70 75 80

Xaa Xaa Xaa Xaa Xaa Val Xaa Leu Xaa Xaa Xaa Xaa Xaa Met Xaa Val
 85 90 95

Xaa Xaa Cys Xaa Cys Xaa
 100

<210> 7
 <211> 97
 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Generic Sequence 9

<220>
<221> MOD_RES
<222> (1)..(97)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (1)..(16)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (18)..(24)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (26)..(26)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (28)..(28)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (30)..(61)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (63)..(63)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (65)..(72)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (74)..(93)
<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (95)..(95)
<223> Xaa is independently selected from a group of one or more

specified amino acids as defined in the specification

<220>

<221> misc_feature

<222> (97)..(97)

<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<400> 7

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
1 5 10 15

Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Gly Xaa Cys Xaa Xaa Xaa
20 25 30

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
35 40 45

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Pro
50 55 60

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
65 70 75 80

Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Cys
85 90 95

Xaa

<210> 8

<211> 102

<212> PRT

<213> Artificial Sequence

<220>

<223> Generic Sequence 10

<220>

<221> MOD_RES

<222> (1)..(102)

<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>

<221> misc_feature

<222> (2)..(21)

<223> Xaa is independently selected from a group of one or more specified amino acids as defined in the specification

<220>

<221> misc_feature
 <222> (23)..(29)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (31)..(31)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (33)..(33)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (35)..(66)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (68)..(68)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (70)..(77)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (79)..(98)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (100)..(100)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<220>
 <221> misc_feature
 <222> (102)..(102)
 <223> Xaa is independently selected from a group of one or more
 specified amino acids as defined in the specification

<400> 8

Cys Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa Xaa
 1 5 10 15

Xaa Xaa Xaa Xaa Xaa Pro Xaa Xaa Xaa Xaa Xaa Xaa Xaa Cys Xaa Gly

```

<210> 9
<211> 5
<212> PRT
<213> Artificial Sequence

<220>
<223> Consensus Sequence

<220>
<221> MOD_RES
<222> (1)..(5)
<223> Xaa is independently selected from a group of one or more
        specified amino acids as defined in the specification

<220>
<221> misc_feature
<222> (2)..(5)
<223> Xaa is independently selected from a group of one or more
        specified amino acids as defined in the specification

<400> 9

Cys Xaa Xaa Xaa Xaa
1           5

```

```
<220>
<221> CDS
<222> (49) .. (1341)
```

ggtgcggggcc cggagcccgcg agccccgggta gcgcgtagag ccggcgcg atg cac gtg 57
Met His Val
1

gac agc cgt acc ctc tgg gcc tgc gag gag gcc tgg ctg gtg ttt gac 729
Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu Val Phe Asp
215 220 225

atc	aca	gcc	acc	agc	aac	cac	tgg	gtg	gtc	aat	ccg	cgg	cac	aac	ctg	777
Ile	Thr	Ala	Thr	Ser	Asn	His	Trp	Val	Val	Asn	Pro	Arg	His	Asn	Leu	
		230					235					240				
ggc	ctg	cag	ctc	tcg	gtg	gag	acg	ctg	gat	ggg	cag	agc	atc	aac	ccc	825
Gly	Leu	Gln	Leu	Ser	Val	Glu	Thr	Leu	Asp	Gly	Gln	Ser	Ile	Asn	Pro	
	245					250					255					
aag	ttg	gcg	ggc	ctg	att	ggg	cgg	cac	ggg	ccc	cag	aac	aag	cag	ccc	873
Lys	Leu	Ala	Gly	Leu	Ile	Gly	Arg	His	Gly	Pro	Gln	Asn	Lys	Gln	Pro	
	260				265					270					275	
ttc	atg	gtg	gct	ttc	ttc	aag	gcc	acg	gag	gtc	cac	ttc	cgc	agc	atc	921
Phe	Met	Val	Ala	Phe	Phe	Lys	Ala	Thr	Glu	Val	His	Phe	Arg	Ser	Ile	
				280					285					290		
cgg	tcc	acg	ggg	agc	aaa	cag	cgc	agc	cag	aac	cgc	tcc	aag	acg	ccc	969
Arg	Ser	Thr	Gly	Ser	Lys	Gln	Arg	Ser	Gln	Asn	Arg	Ser	Lys	Thr	Pro	
			295				300						305			
aag	aac	cag	gaa	gcc	ctg	cgg	atg	gcc	aac	gtg	gca	gag	aac	agc	agc	1017
Lys	Asn	Gln	Glu	Ala	Leu	Arg	Met	Ala	Asn	Val	Ala	Glu	Asn	Ser	Ser	
		310					315					320				
agc	gac	cag	agg	cag	gcc	tgt	aag	aag	cac	gag	ctg	tat	gtc	agc	ttc	1065
Ser	Asp	Gln	Arg	Gln	Ala	Cys	Lys	Lys	His	Glu	Leu	Tyr	Val	Ser	Phe	
	325					330					335					
cga	gac	ctg	ggc	tgg	cag	gac	tgg	atc	atc	gcg	cct	gaa	ggc	tac	gcc	1113
Arg	Asp	Leu	Gly	Trp	Gln	Asp	Trp	Ile	Ile	Ala	Pro	Glu	Gly	Tyr	Ala	
	340				345					350					355	
gcc	tac	tac	tgt	gag	ggg	gag	tgt	gcc	ttc	cct	ctg	aac	tcc	tac	atg	1161
Ala	Tyr	Tyr	Cys	Glu	Gly	Glu	Cys	Ala	Phe	Pro	Leu	Asn	Ser	Tyr	Met	
				360					365					370		
aac	gcc	acc	aac	cac	gcc	atc	gtg	cag	acg	ctg	gtc	cac	ttc	atc	aac	1209
Asn	Ala	Thr	Asn	His	Ala	Ile	Val	Gln	Thr	Leu	Val	His	Phe	Ile	Asn	
			375					380					385			
ccg	gaa	acg	gtg	ccc	aag	ccc	tgc	tgt	gcg	ccc	acg	cag	ctc	aat	gcc	1257
Pro	Glu	Thr	Val	Pro	Lys	Pro	Cys	Cys	Ala	Pro	Thr	Gln	Leu	Asn	Ala	
		390					395					400				
atc	tcc	gtc	ctc	tac	ttc	gat	gac	agc	tcc	aac	gtc	atc	ctg	aag	aaa	1305
Ile	Ser	Val	Leu	Tyr	Phe	Asp	Asp	Ser	Ser	Asn	Val	Ile	Leu	Lys	Lys	
	405					410					415					
tac	aga	aac	atg	gtg	gtc	cgg	gcc	tgt	ggc	tgc	cac	tagctcctcc				1351
Tyr	Arg	Asn	Met	Val	Val	Arg	Ala	Cys	Gly	Cys	His					
	420				425				430							
gagaattcag acccttttggg gccaaagtttt tctggatcct ccattgctcg ccttgggccag																1411
gaaccagcag accaactgcc ttttgtgaga ccttccctc cctatcccca acttttaaagg																1471
tgtgagagta ttaggaaaca tgagcagcat atggcttttg atcagttttt cagtggcagc																1531
atccaatgaa caagatccta caagctgtgc aggcaaaacc tagcaggaaa aaaaaacaac																1591

```
gcataaagaa aaatggccgg gccaggcatc tggctgggaa gtctcagcca tgcacggact 1651
cgtttccaga ggtaattatg agcgcctacc agccaggcca cccagccgtg ggaggaaggg 1711
ggcgtggcaa ggggtgggca cattggtgtc tgtgcgaaag gaaaattgac ccggaagttc 1771
ctgtaataaaa tgtcacaata aaacgaatga atgaaaaaaaa aaaaaaaaaa a 1822
```

```
<210> 11
<211> 431
<212> PRT
<213> Homo sapiens
```

```
<400> 11
```

```
Met His Val Arg Ser Leu Arg Ala Ala Ala Pro His Ser Phe Val Ala
1 5 10 15
```

```
Leu Trp Ala Pro Leu Phe Leu Leu Arg Ser Ala Leu Ala Asp Phe Ser
20 25 30
```

```
Leu Asp Asn Glu Val His Ser Ser Phe Ile His Arg Arg Leu Arg Ser
35 40 45
```

```
Gln Glu Arg Arg Glu Met Gln Arg Glu Ile Leu Ser Ile Leu Gly Leu
50 55 60
```

```
Pro His Arg Pro Arg Pro His Leu Gln Gly Lys His Asn Ser Ala Pro
65 70 75 80
```

```
Met Phe Met Leu Asp Leu Tyr Asn Ala Met Ala Val Glu Glu Gly Gly
85 90 95
```

```
Gly Pro Gly Gly Gln Gly Phe Ser Tyr Pro Tyr Lys Ala Val Phe Ser
100 105 110
```

```
Thr Gln Gly Pro Pro Leu Ala Ser Leu Gln Asp Ser His Phe Leu Thr
115 120 125
```

```
Asp Ala Asp Met Val Met Ser Phe Val Asn Leu Val Glu His Asp Lys
130 135 140
```

```
Glu Phe Phe His Pro Arg Tyr His His Arg Glu Phe Arg Phe Asp Leu
145 150 155 160
```

```
Ser Lys Ile Pro Glu Gly Glu Ala Val Thr Ala Ala Glu Phe Arg Ile
165 170 175
```

Tyr Lys Asp Tyr Ile Arg Glu Arg Phe Asp Asn Glu Thr Phe Arg Ile
 180 185 190

Ser Val Tyr Gln Val Leu Gln Glu His Leu Gly Arg Glu Ser Asp Leu
 195 200 205

Phe Leu Leu Asp Ser Arg Thr Leu Trp Ala Ser Glu Glu Gly Trp Leu
 210 215 220

Val Phe Asp Ile Thr Ala Thr Ser Asn His Trp Val Val Asn Pro Arg
 225 230 235 240

His Asn Leu Gly Leu Gln Leu Ser Val Glu Thr Leu Asp Gly Gln Ser
 245 250 255

Ile Asn Pro Lys Leu Ala Gly Leu Ile Gly Arg His Gly Pro Gln Asn
 260 265 270

Lys Gln Pro Phe Met Val Ala Phe Phe Lys Ala Thr Glu Val His Phe
 275 280 285

Arg Ser Ile Arg Ser Thr Gly Ser Lys Gln Arg Ser Gln Asn Arg Ser
 290 295 300

Lys Thr Pro Lys Asn Gln Glu Ala Leu Arg Met Ala Asn Val Ala Glu
 305 310 315 320

Asn Ser Ser Ser Asp Gln Arg Gln Ala Cys Lys Lys His Glu Leu Tyr
 325 330 335

Val Ser Phe Arg Asp Leu Gly Trp Gln Asp Trp Ile Ile Ala Pro Glu
 340 345 350

Gly Tyr Ala Ala Tyr Tyr Cys Glu Gly Glu Cys Ala Phe Pro Leu Asn
 355 360 365

Ser Tyr Met Asn Ala Thr Asn His Ala Ile Val Gln Thr Leu Val His
 370 375 380

Phe Ile Asn Pro Glu Thr Val Pro Lys Pro Cys Cys Ala Pro Thr Gln
 385 390 395 400

Leu Asn Ala Ile Ser Val Leu Tyr Phe Asp Asp Ser Ser Asn Val Ile
 405 410 415

Leu Lys Lys Tyr Arg Asn Met Val Val Arg Ala Cys Gly Cys His
 420 425 430

<210> 12
 <211> 3151
 <212> DNA
 <213> Mus musculus

<400> 12
 ggaggctggg cggttccgac tgcctgacgc cagcgcgacc agatcaatcc aggctcgggg 60
 agcgagcggg cgcaccaagg cgaggccggg gccgaggcgc ggggacggcg gcccgagact 120
 aagcagagcg cggggacggc ggccgggagc ggatcggagc acgggacccg gcgccgggtc 180
 tcgtgcgtcc ctgcggatgg gcgcgccgcc gagccggcgc taactgggat cctcgctgga 240
 acaggagggg cagtattttc tacctttcca aaccgcagac caagaagcta aggagaatct 300
 atgtaaatat actgaaatct ctgttggctc tgcgcccaac accccggagc tggcacctca 360
 ccctgtctga ggagcgtgta gaactagacc agccgctatg aatgtgacca gcttgthtttc 420
 attcacaagt ccagctgtga agagactcct tgggtggaaa cagggcgatg aagaagagaa 480
 atgggcagag aaagctgtgg acgctttggt gaagaaactg aagaagaaga aaggggccat 540
 ggaagagctg gagaaggccc tgagctgccc tggacagccg agtaactgcg tcaccattcc 600
 tcgctccctg gatggcaggt tgcaggtgtc ccaccggaag ggactacctc atgtcattta 660
 ttgccgtgtg tggcgtggc ccgacctcca gagccaccat gaactgaagc ctctggaatg 720
 ctgtgagttc ccatttggtt ccaagcagaa ggaggtctgc atcaaccctt accactataa 780
 gcgagtggag agcccggttc tcccgcgggt gctgggtccg aggcacagcg agtacaatcc 840
 tcagcacagc cttctggctc agttccgcaa cctgggacaa aatgagcctc acatgccact 900
 gaacgccacg ttcccagact ctttccagca gccaacagc caccggttc cccactcccc 960
 caacagcagc taccccaact ctctggcgg cagcagcagc acctaccctc actccccaac 1020
 cagctcagac ccgggcagcc cttttcagat gccagctgac acacccccac ctgcttacct 1080
 gcctcctgaa gaccccatgg ccaggtatgg ctctcagccc atggacacga acatgatggc 1140
 gcctccactg ccgctgaaa tcagcagagg agatgttcag gcagttgctt acgaggaacc 1200
 aaaacactgg tgctctattg tgtactatga gctcaacaac cgtgtgggtg aagcgttcca 1260
 cgctcgtcc accagcgtgc tgggtggatgg tttcacagat ccgtccaaca ataagaaccg 1320
 cttctgcctt ggcttgctct ccaacgttaa ccggaattcc actattgaaa acaccaggcg 1380
 acatattggg aaaggagtcc acctttatta cgttggagga gaggtgtatg cggaatgcct 1440
 cagtgacagc agcatcttcg tgcagagccg gaactgcaac taccaccacg gctttcacc 1500

```

caccaccgtc tgcaagatcc ccagcgggtg cagcttgaaa atcttcaaca accaagagtt 1560
tgctcagcta ctggcgcagt ctgtgaacca cgggttcgag accgtgtatg aactcaccaa 1620
aatgtgcact attcggatga gcttcgtgaa ggggttgagg gccgaatacc accggcgagg 1680
tgttaccagc accccctgct ggattgagat ccatctgcat ggccctctcc agtggctgga 1740
taaggttctg acccagatgg gctcacccca caatcctatt tcatccgtgt cttaaaagac 1800
ctgtggcttc cgtctcttgc aaactatcga gccttgcatg tacttgaagg atggacaagt 1860
cagacaggat ggagacctga cgaaggagcc acgataatac ttgacctctg tgaccaacta 1920
ttggattgag aaactgacaa gccttggttg atagcaagaa ccctttcagt ttacattgtg 1980
acattctgtt gtaaaaatca actaaaatgc tgactttcag caggactttt gtgtatagtt 2040
aaaaaaaaa gagatggcca agccaggagc aaattatcta ttaggaaaaa agaaaaaaat 2100
gattgtaatc aatccttttg tgtggggtgt tggcagaagg ttggcgctga tagtctttct 2160
gaagtgggct ttcacaggc tcagagccca cgttgaatca tcttctcatg ggttttctta 2220
atattttaaa actacttggt tagaaatgaa tgggtttttt gtttgttttt aaagtacagg 2280
ttaatcgtaa tgacatgcat agtaatcttt ctgaaactgt atgctggctg tattactgtc 2340
agaatgatgg caggcatatg ctctttgcta aatatgtata tacagaatat ttggaggtaa 2400
tgaatagtct aaatggctag tgggtttaca gagtatctga ggggcggggg cgggaagaaa 2460
acgacggctg caaatgtaga ctataccgta aagctcagct tgctgcctta aacagacaag 2520
ctggtgtctg aatttgctgt gtttcagttt ttgtagagtt ttatctgact tcttttcttc 2580
tgtcttatcc gctccacggc acagttaagc agctgggtta ttcctctaac tgtgagagca 2640
gatgagtaat tcttctgtt cgcaaatac ctggcttcgt gtttcagtac ccagtatatg 2700
aaaagcttga attgaatgag cagtttttat ggagtttaca gtacagacat aggctttgat 2760
ttccaaataa attgtttgcc aaacctggta actctgttca ttattcgcag gattaaagat 2820
ctctctattg gaatccatct caaagggtgt tttttttgtt tttgtttttg ttttttgttt 2880
tatttttgatt tgtttttttt tgtactatct gggttctttt cttctgttaa tttttttatt 2940
ctcctttgct cttatacagc gagtactttt attccaacac tagcagggtt tttctctact 3000
ggaaattttt aaataaaacc tgtcattatt gcttactttg attaaaaaaa aaaaaaaaaa 3060
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 3120
aaaaaaaaa aaaatttcat ctgtgtctta a 3151

```

<210> 13
<211> 465

<212> PRT

<213> Mus musculus

<400> 13

Met	Asn	Val	Thr	Ser	Leu	Phe	Ser	Phe	Thr	Ser	Pro	Ala	Val	Lys	Arg
1				5					10					15	

Leu	Leu	Gly	Trp	Lys	Gln	Gly	Asp	Glu	Glu	Glu	Lys	Trp	Ala	Glu	Lys
		20					25						30		

Ala	Val	Asp	Ala	Leu	Val	Lys	Lys	Leu	Lys	Lys	Lys	Lys	Gly	Ala	Met
		35				40						45			

Glu	Glu	Leu	Glu	Lys	Ala	Leu	Ser	Cys	Pro	Gly	Gln	Pro	Ser	Asn	Cys
	50					55					60				

Val	Thr	Ile	Pro	Arg	Ser	Leu	Asp	Gly	Arg	Leu	Gln	Val	Ser	His	Arg
65					70					75					80

Lys	Gly	Leu	Pro	His	Val	Ile	Tyr	Cys	Arg	Val	Trp	Arg	Trp	Pro	Asp
				85					90					95	

Leu	Gln	Ser	His	His	Glu	Leu	Lys	Pro	Leu	Glu	Cys	Cys	Glu	Phe	Pro
			100					105					110		

Phe	Gly	Ser	Lys	Gln	Lys	Glu	Val	Cys	Ile	Asn	Pro	Tyr	His	Tyr	Lys
		115					120					125			

Arg	Val	Glu	Ser	Pro	Val	Leu	Pro	Pro	Val	Leu	Val	Pro	Arg	His	Ser
	130					135						140			

Glu	Tyr	Asn	Pro	Gln	His	Ser	Leu	Leu	Ala	Gln	Phe	Arg	Asn	Leu	Gly
145					150					155					160

Gln	Asn	Glu	Pro	His	Met	Pro	Leu	Asn	Ala	Thr	Phe	Pro	Asp	Ser	Phe
				165					170					175	

Gln	Gln	Pro	Asn	Ser	His	Pro	Phe	Pro	His	Ser	Pro	Asn	Ser	Ser	Tyr
			180					185					190		

Pro	Asn	Ser	Pro	Gly	Gly	Ser	Ser	Ser	Thr	Tyr	Pro	His	Ser	Pro	Thr
		195					200					205			

Ser	Ser	Asp	Pro	Gly	Ser	Pro	Phe	Gln	Met	Pro	Ala	Asp	Thr	Pro	Pro
		210				215					220				

Pro Ala Tyr Leu Pro Pro Glu Asp Pro Met Ala Gln Asp Gly Ser Gln
 225 230 235 240

Pro Met Asp Thr Asn Met Met Ala Pro Pro Leu Pro Ala Glu Ile Ser
 245 250 255

Arg Gly Asp Val Gln Ala Val Ala Tyr Glu Glu Pro Lys His Trp Cys
 260 265 270

Ser Ile Val Tyr Tyr Glu Leu Asn Asn Arg Val Gly Glu Ala Phe His
 275 280 285

Ala Ser Ser Thr Ser Val Leu Val Asp Gly Phe Thr Asp Pro Ser Asn
 290 295 300

Asn Lys Asn Arg Phe Cys Leu Gly Leu Leu Ser Asn Val Asn Arg Asn
 305 310 315 320

Ser Thr Ile Glu Asn Thr Arg Arg His Ile Gly Lys Gly Val His Leu
 325 330 335

Tyr Tyr Val Gly Gly Glu Val Tyr Ala Glu Cys Leu Ser Asp Ser Ser
 340 345 350

Ile Phe Val Gln Ser Arg Asn Cys Asn Tyr His His Gly Phe His Pro
 355 360 365

Thr Thr Val Cys Lys Ile Pro Ser Gly Cys Ser Leu Lys Ile Phe Asn
 370 375 380

Asn Gln Glu Phe Ala Gln Leu Leu Ala Gln Ser Val Asn His Gly Phe
 385 390 395 400

Glu Thr Val Tyr Glu Leu Thr Lys Met Cys Thr Ile Arg Met Ser Phe
 405 410 415

Val Lys Gly Trp Gly Ala Glu Tyr His Arg Gln Asp Val Thr Ser Thr
 420 425 430

Pro Cys Trp Ile Glu Ile His Leu His Gly Pro Leu Gln Trp Leu Asp
 435 440 445

Lys Val Leu Thr Gln Met Gly Ser Pro His Asn Pro Ile Ser Ser Val
 450 455 460

Ser
465

<210> 14
<211> 2449
<212> DNA
<213> Mus musculus

<400> 14
ggttaaaata actatctgag atttgttttg ctggtgttgt tgtttaagga aaattaaggt 60
agtaccatat cttaaatacat tgcaacaaga ggcagtattg ctacttataa aagtaaataa 120
tagtgtataa aattgtgttt caaccgaatc ttactggcat ctttctctct tcttggtgaaa 180
cactccatga aacaatagat gcagtagatc aggatgatga ggacgggaat gggggcacta 240
ctacactact atactactac actctaggat tgcaggctgc atgcagagtt aacaacagtc 300
agctgactgt ttacctgaaa gactggcata gaataggaaa atttggtgcc aagtgcataa 360
aaataagcaa atgaaaagac attaatctct ggttagattta ccgggctttt tctgagtgtg 420
gattgttacc tttggtaaga aaatgtcgtc catcttgcca ttcactccgc cagtgggtgaa 480
gagacttctg ggatggaaaa aatcagccgg tgggtctgga ggagcaggtg gtggagagca 540
gaatggacag gaagaaaagt ggtgtgaaaa agcagtgaaa agtctggtga aaaagctaaa 600
gaaaacagga cggtttagatg agcttgagaa agccatcacc actcagaatt gcaataactaa 660
atgtgtcacc ataccaagca cttgctctga aatttgggga ctgagtacag caaatacgggt 720
agatcagtgg gacacaacag gcctttacag cttctctgaa caaaccaggt ctcttgatgg 780
ccgtcttcag gtttcacacc ggaaagggtt gccacatgtt atatatgcc ggctctggcg 840
ctggccggac cttcacagtc atcatgagct caaggcaatc gaaaactgcg aatatgcttt 900
taatctgaaa aaagatgaag tgtgtgtaaa tccgtaccac taccagagag ttgagacccc 960
agtcttgctt ccagtcttag tgccctcggca cacggagatt ctaacagaac tgccaccctt 1020
ggatgactac acccactcca ttccagaaaa cacaaatttc ccagcaggaa ttgagccaca 1080
gagtaattac atcccagaaa caccaccacc tggatatatc agtgaagatg gagaaacaag 1140
tgaccaacag ttgaaccaa gtatggacac aggcctctccg gctgaactgt ctctactac 1200
tctctctcct gttaatcaca gcttggattt gcagccagtt acttactcgg aacctgcatt 1260
ctggtgttca atcgcatact atgaactaaa ccagagggtt ggagagacct tccatgcgtc 1320
acagccctcg ctcaactgtag acggcttcac agaccocatc aactcggaga gggtctgctt 1380
aggcttgctc tccaacgtta accgaaatgc cactgtagaa atgacaagaa gacatatagg 1440
aaggggagtg cgcttgtatt acataggtgg ggaagtgttt gctgagtgcc taagtgatag 1500

```

tgcaatcttt gtgcagagcc ccaactgtaa ccagagatac ggctggcacc ctgcaacagt 1560
gtgtaagatc ccaccaggct gtaacctgaa gatcttcaac aaccaagaat ttgctgctct 1620
tctggctcag tctgtcaacc agggttttga agccgtttat cagctaaccc gaatgtgcac 1680
cataagaatg agttttgtga agggctgggg agcagaatat cggaggcaga cagtaacaag 1740
tactccttgc tggattgaac ttcatctgaa tggccctctg cagtggctgg acaaagtatt 1800
aactcagatg ggatcccctt cagtgcgatg ctcaagcatg tcgtaaacc atcaaagact 1860
cgctgtaaca gtcctccgt cgtagtattc atgtatgac ccgtggactg tttgctatcc 1920
aaaaattcca gagcaaaaac agcacttgag gtctcatcag ttaaagcacc ttgtggaatc 1980
tgtttcctat atttgaatat tagatgggaa aattagtgtc tagaaatgcc ctcccagcg 2040
gggaaaaaga agacttaaag acttaatgat gtcttggttg gcataagaca gtatcccaaa 2100
ggttattaat aacagtagta gttgtgtaca ggtaatgtgt ccagaccagcag tattgcagta 2160
ctatgctgtt tgtatacatt cttagtttgc ataaatgagg tgtgtgtgct gcttcttggt 2220
ctaggcaagc ctttataaaa ttacagtatc taatctgtta ttcccacttc tccgttattt 2280
ttgtgtcttt tttaatatat aatatatata tatcaagatt ttcaaattat catttagaag 2340
cagattttcc ttgtagaaac taatttttct gccttttacc aaaaataaac aaactcttgg 2400
gggaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 2449

```

```

<210> 15
<211> 494
<212> PRT
<213> Mus musculus

```

```

<400> 15

```

```

Met Ser Ser Ile Leu Pro Phe Thr Pro Pro Val Val Lys Arg Leu Leu
1           5           10           15

```

```

Gly Trp Lys Lys Ser Ala Gly Gly Ser Gly Gly Ala Gly Gly Gly Glu
          20           25           30

```

```

Gln Asn Gly Gln Glu Glu Lys Trp Cys Glu Lys Ala Val Lys Ser Leu
          35           40           45

```

```

Val Lys Lys Leu Lys Lys Thr Gly Arg Leu Asp Glu Leu Glu Lys Ala
          50           55           60

```

```

Ile Thr Thr Gln Asn Cys Asn Thr Lys Cys Val Thr Ile Pro Ser Thr
65           70           75           80

```

Cys Ser Glu Ile Trp Gly Leu Ser Thr Ala Asn Thr Val Asp Gln Trp
 85 90 95

Asp Thr Thr Gly Leu Tyr Ser Phe Ser Glu Gln Thr Arg Ser Leu Asp
 100 105 110

Gly Arg Leu Gln Val Ser His Arg Lys Gly Leu Pro His Val Ile Tyr
 115 120 125

Cys Arg Leu Trp Arg Trp Pro Asp Leu His Ser His His Glu Leu Lys
 130 135 140

Ala Ile Glu Asn Cys Glu Tyr Ala Phe Asn Leu Lys Lys Asp Glu Val
 145 150 155 160

Cys Val Asn Pro Tyr His Tyr Gln Arg Val Glu Thr Pro Val Leu Pro
 165 170 175

Pro Val Leu Val Pro Arg His Thr Glu Ile Leu Thr Glu Leu Pro Pro
 180 185 190

Leu Asp Asp Tyr Thr His Ser Ile Pro Glu Asn Thr Asn Phe Pro Ala
 195 200 205

Gly Ile Glu Pro Gln Ser Asn Tyr Ile Pro Glu Thr Pro Pro Pro Gly
 210 215 220

Tyr Ile Ser Glu Asp Gly Glu Thr Ser Asp Gln Gln Leu Asn Gln Ser
 225 230 235 240

Met Asp Thr Gly Ser Pro Ala Glu Leu Ser Pro Thr Thr Leu Ser Pro
 245 250 255

Val Asn His Ser Leu Asp Leu Gln Pro Val Thr Tyr Ser Glu Pro Ala
 260 265 270

Phe Trp Cys Ser Ile Ala Tyr Tyr Glu Leu Asn Gln Arg Val Gly Glu
 275 280 285

Thr Phe His Ala Ser Gln Pro Ser Leu Thr Val Asp Gly Phe Thr Asp
 290 295 300

Pro Ser Asn Ser Glu Arg Phe Cys Leu Gly Leu Leu Ser Asn Val Asn
 305 310 315 320

Arg Asn Ala Thr Val Glu Met Thr Arg Arg His Ile Gly Arg Gly Val
 325 330 335
 Arg Leu Tyr Tyr Ile Gly Gly Glu Val Phe Ala Glu Cys Leu Ser Asp
 340 345 350
 Ser Ala Ile Phe Val Gln Ser Pro Asn Cys Asn Gln Arg Tyr Gly Trp
 355 360 365
 His Pro Ala Thr Val Cys Lys Ile Pro Pro Gly Cys Asn Leu Lys Ile
 370 375 380
 Phe Asn Asn Gln Glu Phe Ala Ala Leu Leu Ala Gln Ser Val Asn Gln
 385 390 395 400
 Gly Phe Glu Ala Val Tyr Gln Leu Thr Arg Met Cys Thr Ile Arg Met
 405 410 415
 Ser Phe Val Lys Gly Trp Gly Ala Glu Tyr Arg Arg Gln Thr Val Thr
 420 425 430
 Ser Thr Pro Cys Trp Ile Glu Leu His Leu Asn Gly Pro Leu Gln Trp
 435 440 445
 Leu Asp Lys Val Leu Thr Gln Met Gly Ser Pro Ser Val Arg Cys Ser
 450 455 460
 Ser Met Ser Glu Ile Thr Ala Cys Ile Ser Gln Arg Gln Val Lys Glu
 465 470 475 480
 Leu Asp Val Leu Ile Gln Thr Glu Gln Trp Pro Gln Tyr Pro
 485 490

<210> 16
 <211> 6256
 <212> DNA
 <213> Homo sapiens

<400> 16
 gcggccgccc cctccgcccc gcgttcgggg ccttcccgac cctgcactgc tgccgtccgc 60
 ccgcccggcc gctctttctct tcgccgtggg agccgctccg ggcgaggggc cgcgcgccga 120
 gccccgcagg ctgcagcgcc gcggcccggc ccggcgcccc ggcaacttcg ccgagagttg 180
 aggcgaagtt tgggcgaccg cggcaggccc cgcccgagct cccctctgcg cccccggcgt 240
 cccgtcgagc ccagccccgc cggggggcgt cctcgccgcc cgcgcgccct cccagccat 300

gtcgtccatc	ctgcctttca	ctcccccgat	cgtgaagcgc	ctgctgggct	ggaagaaggg	360
cgagcagaac	gggcaggagg	agaaatgggtg	cgagaaggcg	gtcaagagcc	tgggtcaagaa	420
actcaagaag	acgggggcagc	tggacgagct	ggagaaggcc	atcaccacgc	agaacgtcaa	480
caccaagtgc	atcaccatcc	ccagggtccct	ggatggcccg	ttgcagggtgt	cccatcgga	540
ggggctccct	catgtcatct	actgccgcct	gtggcgatgg	ccagacctgc	acagccacca	600
cgagctacgg	gccatggagc	tgtgtgagtt	cgccttcaat	atgaagaagg	acgaggtctg	660
cgtgaatccc	taccactacc	agagagtaga	gacaccagtt	ctacctcctg	tgttggtgcc	720
acgccacaca	gagatcccg	ccgagttccc	cccactggac	gactacagcc	attccatccc	780
cgaaaacact	aacttccccg	caggcatcga	gccccagagc	aatattccag	agaccccacc	840
ccctggctac	ctgagtgaag	atggagaaac	cagtgaccac	cagatgaacc	acagcatgga	900
cgcaggttct	ccaaacctat	ccccgaatcc	gatgtcccca	gcacataata	acttggaacct	960
gcagccagtt	acctactgcg	agccggcctt	ctgggtgctcc	atctcctact	acgagctgaa	1020
ccagcgcgtc	ggggagacat	tccacgcctc	gcagccatcc	atgactgtgg	atggcttcac	1080
cgacccctcc	aattcggagc	gcttctgcct	agggctgctc	tccaatgtca	acaggaatgc	1140
agcagtggag	ctgacacgga	gacacatcgg	aagaggcgtg	cggctctact	acatcggagg	1200
ggaggtcttc	gcagagtgcc	tcagtgcag	cgctatTTTT	gtccagtctc	ccaactgtaa	1260
ccagcgtat	ggctggcacc	cggccaccgt	ctgcaagatc	ccaccaggat	gcaacctgaa	1320
gatcttcaac	aaccaggagt	tcgctgcct	cctggcccag	tcgggtcaacc	agggtcttga	1380
ggctgtctac	cagttgacct	gaatgtgcac	catccgcatg	agcttcgtca	aaggctgggg	1440
agcggagtac	aggagacaga	ctgtgaccag	tacccccctgc	tggattgagc	tgcacctgaa	1500
tgggcctttg	cagtggcttg	acaaggtcct	caccagatg	ggctcccaa	gcatccgctg	1560
ttccagtgtg	tcttagagac	atcaagtatg	gtaggggagg	gcaggcttgg	ggaaaatggc	1620
catgcaggag	gtggagaaaa	ttggaactct	actcaacca	ttgttgtaa	ggaagaagaa	1680
atctttctcc	ctcaactgaa	ggggtgcacc	cacctgtttt	ctgaaacaca	cgagcaaacc	1740
cagaggtgga	tgttatgaac	agctgtgtct	gccaacaca	tttacccttt	ggccccactt	1800
tgaagggcaa	gaaatggcgt	ctgctctggg	ggcttaagtg	agcagaacag	gtagtattac	1860
accaccggcc	ccctcccccc	agactctttt	tttgagtgc	agctttctgg	gatgtcacag	1920
tccaaccaga	aacaccctc	tgtctaggac	tgcagtgtgg	agttcacctt	ggaagggcgt	1980
tctaggtagg	aagagcccg	agggccatgc	agacctcatg	cccagctctc	tgacgcttgt	2040
gacagtgcct	cttccagtga	acattcccag	cccagccccg	ccccgccccg	ccccaccact	2100

ccagcagacc	ttgcccccttg	tgagctggat	agacttggga	tggggaggga	gggagttttg	2160
tctgtctccc	tccccctctca	gaacatactg	attgggaggt	gcgtgttcag	cagaacctgc	2220
acacaggaca	gcgggaaaaa	tcgatgagcg	ccacctcttt	aaaaactcac	ttacgtttgt	2280
cctttttcac	tttgaaaagt	tggaaggatc	tgctgaggcc	cagtgcataat	gcaatgtata	2340
gtgtctatta	tcacattaat	ctcaaagaga	ttcgaatgac	ggtaagtgtt	ctcatgaagc	2400
aggaggccct	tgtcgtggga	tggcatttgg	tctcaggcag	caccacactg	ggtgcgtctc	2460
cagtcatctg	taagagcttg	ctccagattc	tgatgcatac	ggctatatattg	gtttatgtag	2520
tcagttgcat	tcattaaatc	aactttatca	tatgctcttt	taaatgtttg	gtttatatat	2580
tttctttaaa	aatcctgggc	tggcacattg	actgggaaac	ctgagtgaga	cccagcaact	2640
gcttctctcc	cttctctctc	ctgaggtgaa	gcttttccag	gttttgttga	agagatacct	2700
gccagcactt	ctgcaagctg	aaattttacag	aagcaaattc	accagaaggg	aaacatctca	2760
ggccaacata	ggcaaataa	aagggtctatt	aaaatatatt	tacacctttg	aaaattgcag	2820
gcttggtaca	aagaggtctg	tcattctccc	cctgggatata	aagatgatct	agctcccggg	2880
agaggatcac	cggtgacaac	tatagcagtt	gtattgtgta	acaagtactg	ctcccagcag	2940
caattaggga	gaaaactagt	ctaaattatt	tcaactggaa	aaaagaaaaa	agagtcctct	3000
tctttttcca	gccttttgca	gaacacagta	gacagaactg	ccaccttcaa	ttggtacttt	3060
attctttgct	gctgtttttg	tataaaatga	cctatcccac	gtttttgcat	gaatttatag	3120
caggaaaaat	caagggtatt	cctatggaag	tcctgcttta	ttccaggtga	agggaaggaa	3180
gtgtatatac	ttttggcaag	tcatacagct	caaattgtgat	gagatttctg	atgttagagg	3240
gagatggaga	ggcttcctga	tgctctatct	gcagggtcct	gtgcctctga	agttctagcc	3300
atgaggtttc	caggtaggac	agctgctccc	caagcctcct	gaggacacag	gaagagacgg	3360
aaggagcacc	ttgacagact	tgtgtgagtc	ttctcgaagg	agggttgact	cagaacctcag	3420
agacaataca	aaacctctca	cttcctctga	gagggccaaa	tgctgtgagt	ctgaagtatg	3480
tgcttgggtg	gaaatgatct	atggcctggt	tcttacacag	gaagccccct	gaacctcctg	3540
tacatgtgtt	catgtttcca	gccagctctg	agaccagga	accaaataat	ccattttggc	3600
ttctgctaga	gcagtcatgg	ttcctctcct	aaaagccatg	ggcagcagtt	tccgaggggc	3660
tgcatgatcc	acctgctgca	cgatcctatg	agggttcct	gtggcacaca	gccctctggg	3720
tgcttgggaa	ctagcttcag	gcacagcctg	attctggtga	tccagtgatc	tatggaagtc	3780
gtgtcttact	ccaggtgaag	ggggaaaaaa	aaagcctata	ctttggcagg	ttatgaactt	3840
tgaatgtgat	gaaatgacac	gtttggctgc	atttggatgg	tgtcttagaa	ccctcattgc	3900

tcagacctga	aggctacttc	taggagcatg	aagtttgagt	tttgtgtttt	tccaaaggat	3960
acttccttgg	ccctttttct	ttattgacta	gaccaccaga	ggaggatgtg	tgggattgta	4020
ggcaaaccga	cctgtggcat	cactgaaaat	aaatttgatc	atacctaaga	ggttaggaaa	4080
tggtgccatt	cccaccttag	agtgctacat	aggtgctttg	ggcgtatgta	acattagtgt	4140
ccttccttga	agccacaagc	tagttttctt	agttttaaaa	tcctgttgta	tgaatggcat	4200
ttgtatatta	aaacactttt	ttaaaggaca	gttgaaaagg	gcaagaggaa	accagggcag	4260
ttctagagga	gtgctgggtga	ctggatagca	gttttaagtg	gcgttcacct	agtcaacacg	4320
accgcgtgtg	ttgccccctg	cctgggctcc	ccgccatgac	atcttcacct	tgcagcttgt	4380
gctgagactg	acccaagtgc	agctagcact	gggacacaga	tccttgtctt	cagcaccttc	4440
caaggagcca	actttttatt	ccttttcctc	ctccccctcc	cacctcgctt	cttcccaatt	4500
tagtaactta	gatgcttcca	gcacatacgt	aggtagctac	cccagccggg	ttggattaca	4560
ggcctgtgct	ggaacatcat	ctcagttggc	caccttcctg	gcaggctgta	gacctgacat	4620
tttgagacaa	gcctagagtc	aggagcaggg	actttgactc	ttaggaagag	cacacatgag	4680
ggcaaggctg	ctggcagacg	tctccattgt	ccttatgttg	tctgtgttgt	atTTTTTTTT	4740
ttttattgac	catggtgatt	atTTTTTTaa	accatcgttt	atatactgaa	gtgagctata	4800
gcacatatca	tgtgcttagt	ttgtttattt	ttctccatct	ccccttggct	tcctagagtt	4860
tggacatatt	ccaggctaaa	tgcttttact	caagactaca	gaaaggtttg	aagtagtgtg	4920
tgcattggcat	gcacgtatgt	aagtaatctg	gggaagaagc	aaagatctgt	ttcattctta	4980
gcctcaggcc	tcattgagggt	ctccacaggg	ccggagctca	ggttacacca	ctccttcgtc	5040
cttacaggag	atgtagggag	aagaatctgc	aggctgcttg	taggactgtt	caccaagggg	5100
gataccagca	gcaagagagt	gcacccgttt	agccctggac	cctgtttctt	actgtgtgac	5160
ttggctagag	ttgggagttc	ccccaaaata	aacgtgtccc	cattttacca	gaaccaaacc	5220
tcaacacagc	gaagctgtac	tgtctttgtg	tggcaaagat	gttcccttgt	aggccccctt	5280
caggtaaccg	tcttcacaat	gtattttcat	cacagtttaa	ggagcatcag	ccgcttctca	5340
agtgggtagg	gaaagcagaa	aaacgtacgc	aagaggacat	ggatccaaaa	tgatgatgaa	5400
gcatctccca	tggggaggtg	atggtgggga	gatgatgggc	taaacaggca	acttttcaaa	5460
aacacagcta	tcataaaaa	gaaacttgcc	tcattgtaaac	tggattgaga	aattctcagt	5520
gattctgcaa	tggatttttt	tttaattgcag	aagtaatgta	tactctagta	ttctgggtgt	5580
tttatattta	tgtataaatt	tcttaaaacc	attcagacag	ataactattt	aatttttttt	5640
aagaaagtgt	gaaaggctct	tcctcccaag	gacagtggct	ggaagagtgt	gggcacagcc	5700

```

agttctgaat gttggtggag ggtgtagtgg ctttttggct cagcatccag aaacaccaaa 5760
ccaggctggc taaacaagtg gccgcgtgta aaaacagaca gctctgagtc aaatctgggc 5820
ccttcacaaa gggtcctctg aaccaagccc cactcccttg ctaggggtga aagcattaca 5880
gagagatgga gccatctatc caagaagcct tcactcacct tcactgctgc tgttgcaact 5940
cggctgttct ggactctgat gtgtgtggag ggatggggaa tagaacattg actgtgttga 6000
ttaccttcac tattcggcca gcctgacctt ttaataactt tgtaaaaagc atgtatgtat 6060
ttatagtgtt ttagattttt ctaactttta tatcttaaaa gcagagcacc tgtttaagca 6120
ttgtaccctt attgttaaag atttgtgtcc tctcattccc tctcttcctc ttgtaagtgc 6180
ccttctaata aacttttcat ggaaaagctc ctgtgccagg agctcagtct gaaaaaaaaa 6240
aaaaaaaaaa aaaaaa 6256

```

```

<210> 17
<211> 425
<212> PRT
<213> Homo sapiens

```

```

<400> 17

```

```

Met Ser Ser Ile Leu Pro Phe Thr Pro Pro Ile Val Lys Arg Leu Leu
1           5           10           15

```

```

Gly Trp Lys Lys Gly Glu Gln Asn Gly Gln Glu Glu Lys Trp Cys Glu
20           25           30

```

```

Lys Ala Val Lys Ser Leu Val Lys Lys Leu Lys Lys Thr Gly Gln Leu
35           40           45

```

```

Asp Glu Leu Glu Lys Ala Ile Thr Thr Gln Asn Val Asn Thr Lys Cys
50           55           60

```

```

Ile Thr Ile Pro Arg Ser Leu Asp Gly Arg Leu Gln Val Ser His Arg
65           70           75           80

```

```

Lys Gly Leu Pro His Val Ile Tyr Cys Arg Leu Trp Arg Trp Pro Asp
85           90           95

```

```

Leu His Ser His His Glu Leu Arg Ala Met Glu Leu Cys Glu Phe Ala
100          105          110

```

```

Phe Asn Met Lys Lys Asp Glu Val Cys Val Asn Pro Tyr His Tyr Gln
115          120          125

```

Arg Val Glu Thr Pro Val Leu Pro Pro Val Leu Val Pro Arg His Thr
 130 135 140

Glu Ile Pro Ala Glu Phe Pro Pro Leu Asp Asp Tyr Ser His Ser Ile
 145 150 155 160

Pro Glu Asn Thr Asn Phe Pro Ala Gly Ile Glu Pro Gln Ser Asn Ile
 165 170 175

Pro Glu Thr Pro Pro Pro Gly Tyr Leu Ser Glu Asp Gly Glu Thr Ser
 180 185 190

Asp His Gln Met Asn His Ser Met Asp Ala Gly Ser Pro Asn Leu Ser
 195 200 205

Pro Asn Pro Met Ser Pro Ala His Asn Asn Leu Asp Leu Gln Pro Val
 210 215 220

Thr Tyr Cys Glu Pro Ala Phe Trp Cys Ser Ile Ser Tyr Tyr Glu Leu
 225 230 235 240

Asn Gln Arg Val Gly Glu Thr Phe His Ala Ser Gln Pro Ser Met Thr
 245 250 255

Val Asp Gly Phe Thr Asp Pro Ser Asn Ser Glu Arg Phe Cys Leu Gly
 260 265 270

Leu Leu Ser Asn Val Asn Arg Asn Ala Ala Val Glu Leu Thr Arg Arg
 275 280 285

His Ile Gly Arg Gly Val Arg Leu Tyr Tyr Ile Gly Gly Glu Val Phe
 290 295 300

Ala Glu Cys Leu Ser Asp Ser Ala Ile Phe Val Gln Ser Pro Asn Cys
 305 310 315 320

Asn Gln Arg Tyr Gly Trp His Pro Ala Thr Val Cys Lys Ile Pro Pro
 325 330 335

Gly Cys Asn Leu Lys Ile Phe Asn Asn Gln Glu Phe Ala Ala Leu Leu
 340 345 350

Ala Gln Ser Val Asn Gln Gly Phe Glu Ala Val Tyr Gln Leu Thr Arg
 355 360 365

Met Cys Thr Ile Arg Met Ser Phe Val Lys Gly Trp Gly Ala Glu Tyr
 370 375 380

Arg Arg Gln Thr Val Thr Ser Thr Pro Cys Trp Ile Glu Leu His Leu
 385 390 395 400

Asn Gly Pro Leu Gln Trp Leu Asp Lys Val Leu Thr Gln Met Gly Ser
 405 410 415

Pro Ser Ile Arg Cys Ser Ser Val Ser
 420 425

<210> 18
 <211> 3221
 <212> DNA
 <213> Homo sapiens

<400> 18
 caacaacacg gccctggctg tcgtcgccgc tgcggtaacg gagcggtttg ggtggcggag 60
 cctgcgttcg cgccttccca ctcccctcgc caccgcccga gcccagggtta tcctgaatac 120
 atgtctaaca attttccttg caacgtagc tgttgttttt cactgtttcc aaaggatcaa 180
 aattgcttca gaaattggag acatatattga tttaaaagga aaaacttgaa caaatggaca 240
 atatgtctat tacgaataca ccaacaagta atgatgcctg tctgagcatt gtgcatagtt 300
 tgatgtgcca tagacaaggt ggagagagtg aaacatttgc aaaaagagca attgaaagtt 360
 tggtaaagaa gctgaaggag aaaaaagatg aattggattc ttttaataaca gctataacta 420
 caaatggagc tcatcctagt aaatgtgtta ccatacagag aacattggat gggagggttc 480
 aggtggctgg tcggaagga tttcctcatg tgatctatgc ccgtctctgg aggtggcctg 540
 atcttcacaa aaatgaacta aaacatgtta aatattgtca gtatgcgttt gacttaaaat 600
 gtgatagtgt ctgtgtgaat ccatatcact acgaacgagt tgtatcacct ggaattgatc 660
 tctcaggatt aacactgcag agtaatgctc catcaagtat gatggtgaag gatgaatatg 720
 tgcatgactt tgagggacag ccatcgttgt ccaactgaagg acattcaatt caaaccatcc 780
 agcatccacc aagtaatcgt gcatcgacag agacatacag caccacagct ctggttagccc 840
 catctgagtc taatgctacc agcactgcca actttcccaa cattcctgtg gcttccacaa 900
 gtcagcctgc cagtatactg gggggcagcc atagtgaagg actggtgcag atagcatcag 960
 ggcctcagcc aggacagcag cagaatggat ttactgggtca gccagctact taccatcata 1020
 acagcactac cacctggact ggaagtagga ctgcaccata cacacctaatt ttgcctcacc 1080
 accaaaacgg ccattcttcag caccacccgc ctatgccgcc ccatcccgga cattactggc 1140

ctgttcacaa	tgagcttgca	ttccagcctc	ccattttccaa	tcatacctgct	cctgagtatt	1200
ggtgttccat	tgcttacttt	gaaatggatg	ttcaggtagg	agagacattt	aagggttcctt	1260
caagctgccc	tattgttact	gttgatggat	acgtggaccc	ttctggagga	gatcgctttt	1320
gtttgggtca	actctccaat	gtccacagga	cagaagccat	tgagagagca	aggttgcaca	1380
taggcaaagg	tgtgcagttg	gaatgtaaag	gtgaagggtga	tgtttgggtc	aggtgcctta	1440
gtgaccacgc	ggtcttttga	cagagttact	acttagacag	agaagctggg	cgtgcacctg	1500
gagatgctgt	tcataagatc	tacccaagtg	catatataaa	ggtctttgat	ttgcgtcagt	1560
gtcatcgaca	gatgcagcag	caggcggcta	ctgcacaagc	tgacagcagct	gcccaggcag	1620
cagccgtggc	aggaaacatc	cctggcccag	gatcagtagg	tggaatagct	ccagctatca	1680
gtctgtcagc	tgctgctgga	attgggtgtg	atgaccttcg	tcgcttatgc	atactcagga	1740
tgagttttgt	gaaaggctgg	ggaccggatt	acccaagaca	gagcatcaaa	gaaacacctt	1800
gctggattga	aattcactta	caccgggccc	tccagctcct	agacgaagta	cttcatacca	1860
tgccgattgc	agaccacaaa	cctttagact	gaggtctttt	accgttgggg	cccttaacct	1920
tatcaggatg	gtggactaca	aaatacaatc	ctgtttataa	tctgaagata	tattttcactt	1980
ttgttctgct	ttatcttttc	ataaagggtt	gaaaatgtgt	ttgctgcctt	gctcctagca	2040
gacagaaaact	ggattaaaaac	aatttttttt	ttcctcttca	gaacttgtca	ggcatggctc	2100
agagcttgaa	gattaggaga	aacacattct	tattaattct	tcacctgtta	tgtatgaagg	2160
aatcattcca	gtgctagaaa	atttagccct	ttaaaacgtc	ttagagcctt	ttatctgcag	2220
aacatcgata	tgtatatcat	tctacagaat	aatccagtat	tgctgatttt	aaaggcagag	2280
aagttctcaa	agttaattca	cctatgttat	tttgtgtaca	agttgttatt	gttgaacata	2340
cttcaaaaat	aatgtgccat	gtgggtgagt	taattttacc	aagagtaact	ttactctgtg	2400
tttaaaaagt	aagttaataa	tgtattgtaa	tctttcatcc	aaaatatatt	ttgcaagtta	2460
tattagtga	gatggtttca	attcagattg	tcttgcaact	tcagttttat	ttttgccaag	2520
gcaaaaaaact	cttaatctgt	gtgtatatattg	agaatccctt	aaaattacca	gacaaaaaaa	2580
tttaaaatta	cgtttgttat	tcctagtggga	tgactgttga	tgaagtatac	ttttcccttg	2640
ttaaacagta	gttgtattct	tctgtatttc	taggcacaag	gttggttgct	aagaagccta	2700
taagaggaat	ttcttttcct	tcattcatag	ggaaagggtt	tgtatttttt	aaaacactaa	2760
aagcagcgtc	actctaccta	atgtctcact	gttctgcaaa	ggtggcaatg	cttaaaactaa	2820
ataatgaata	aactgaatat	tttggaaaact	gctaaattct	atgttaaata	ctgtgcagaa	2880
taatggaaac	attacagttc	ataataggta	gtttggatat	ttttgtactt	gatttgatgt	2940

```

gactttttttt ggtataatgt ttaaatacatg tatgttatga tattgtttta aattcagttt 3000
ttgtatcttg gggcaagact gcaaactttt ttatatcttt tggttattct aagccctttg 3060
ccatcaatga tcatatcaat tggcagtgac tttgtataga gaatttaagt agaaaagttg 3120
cagatgtatt gactgtacca cagacacaat atgtatgctt tttacctagc tggtagcata 3180
aataaaaactg aatctcaaca taaaaaaaaa aaaaaaaaaa a 3221

```

```

<210> 19
<211> 552
<212> PRT
<213> Homo sapiens

```

```

<400> 19

```

```

Met Asp Asn Met Ser Ile Thr Asn Thr Pro Thr Ser Asn Asp Ala Cys
1          5          10          15

```

```

Leu Ser Ile Val His Ser Leu Met Cys His Arg Gln Gly Gly Glu Ser
          20          25          30

```

```

Glu Thr Phe Ala Lys Arg Ala Ile Glu Ser Leu Val Lys Lys Leu Lys
          35          40          45

```

```

Glu Lys Lys Asp Glu Leu Asp Ser Leu Ile Thr Ala Ile Thr Thr Asn
50          55          60

```

```

Gly Ala His Pro Ser Lys Cys Val Thr Ile Gln Arg Thr Leu Asp Gly
65          70          75          80

```

```

Arg Leu Gln Val Ala Gly Arg Lys Gly Phe Pro His Val Ile Tyr Ala
          85          90          95

```

```

Arg Leu Trp Arg Trp Pro Asp Leu His Lys Asn Glu Leu Lys His Val
100          105          110

```

```

Lys Tyr Cys Gln Tyr Ala Phe Asp Leu Lys Cys Asp Ser Val Cys Val
115          120          125

```

```

Asn Pro Tyr His Tyr Glu Arg Val Val Ser Pro Gly Ile Asp Leu Ser
130          135          140

```

```

Gly Leu Thr Leu Gln Ser Asn Ala Pro Ser Ser Met Met Val Lys Asp
145          150          155          160

```

```

Glu Tyr Val His Asp Phe Glu Gly Gln Pro Ser Leu Ser Thr Glu Gly

```

					165				170						175
His	Ser	Ile	Gln	Thr	Ile	Gln	His	Pro	Pro	Ser	Asn	Arg	Ala	Ser	Thr
			180					185					190		
Glu	Thr	Tyr	Ser	Thr	Pro	Ala	Leu	Leu	Ala	Pro	Ser	Glu	Ser	Asn	Ala
		195					200					205			
Thr	Ser	Thr	Ala	Asn	Phe	Pro	Asn	Ile	Pro	Val	Ala	Ser	Thr	Ser	Gln
	210					215					220				
Pro	Ala	Ser	Ile	Leu	Gly	Gly	Ser	His	Ser	Glu	Gly	Leu	Leu	Gln	Ile
225					230					235					240
Ala	Ser	Gly	Pro	Gln	Pro	Gly	Gln	Gln	Gln	Asn	Gly	Phe	Thr	Gly	Gln
				245					250					255	
Pro	Ala	Thr	Tyr	His	His	Asn	Ser	Thr	Thr	Thr	Trp	Thr	Gly	Ser	Arg
			260					265					270		
Thr	Ala	Pro	Tyr	Thr	Pro	Asn	Leu	Pro	His	His	Gln	Asn	Gly	His	Leu
		275					280					285			
Gln	His	His	Pro	Pro	Met	Pro	Pro	His	Pro	Gly	His	Tyr	Trp	Pro	Val
	290					295					300				
His	Asn	Glu	Leu	Ala	Phe	Gln	Pro	Pro	Ile	Ser	Asn	His	Pro	Ala	Pro
305					310					315					320
Glu	Tyr	Trp	Cys	Ser	Ile	Ala	Tyr	Phe	Glu	Met	Asp	Val	Gln	Val	Gly
				325					330					335	
Glu	Thr	Phe	Lys	Val	Pro	Ser	Ser	Cys	Pro	Ile	Val	Thr	Val	Asp	Gly
			340					345					350		
Tyr	Val	Asp	Pro	Ser	Gly	Gly	Asp	Arg	Phe	Cys	Leu	Gly	Gln	Leu	Ser
		355					360					365			
Asn	Val	His	Arg	Thr	Glu	Ala	Ile	Glu	Arg	Ala	Arg	Leu	His	Ile	Gly
	370					375					380				
Lys	Gly	Val	Gln	Leu	Glu	Cys	Lys	Gly	Glu	Gly	Asp	Val	Trp	Val	Arg
385					390					395					400
Cys	Leu	Ser	Asp	His	Ala	Val	Phe	Val	Gln	Ser	Tyr	Tyr	Leu	Asp	Arg

```
<210> 20
<211> 6504
<212> DNA
<213> Mus musculus
```

<400>	20						
gagagcgcgc	ctgggcggcg	gggaggactt	gcactaagaa	gaagcctatg	gcacctgtca		60
agttaaattgt	cactccccgc	ctccacttgg	actttctgct	taagacctgc	atgtgacttt		120
tcacctgcga	gccacgcttt	tggtatctac	tgactttgat	tacaggaaag	tgtctgaaga		180
tttgtatcaa	atgacgtcaa	tggccagctt	gttttctttc	actagtccag	ccgtgaagcg		240
attgttgggc	tggaaacaag	gtgacgagga	agagaaatgg	gcagaaaagg	cagtggatgc		300
tttagtgaaa	aagctgaaga	agaagaaggg	tgctatggag	gagctggaga	aagccttgag		360
cagcccagga	cagccaagca	agtgtgtcac	gatccccagg	tccttggatg	gacgtctgca		420
agtttctcac	aggaaaggct	tgccccatgt	tatatattgc	cgtgtttggc	gctggccaga		480

tttgcagagc catcacgagc taaaaccatt ggatatttgt gaatttcctt ttggatctaa	540
gcaaaaggaa gtttgtatca atccatacca ctataagaga gtggagagtc cagtcttacc	600
tccagtatta gtgcctcgtc acaatgaatt caatccacaa cacagccttc tggttcagtt	660
caggaacctg agccacaatg aaccgcacat gccacaaaac gccacgtttc ccgattcttt	720
ccaccaaccc aacaacgctc ctttccccctt atctcctaac agcccctatc ctccctcccc	780
tgctagcagc acatatccca actccccagc aagctctgga cctggaagtc catttcaact	840
cccagctgac acccctcccc ctgcctatat gccacctgat gatcagatgg cccagataa	900
ttcccagcct atggatacaa gcagtaacat gattcctcag accatgcca gcatatccag	960
cagagatggt cagcctgtcg cctatgagga gcccaaacac tgggtgttca ttgtctacta	1020
tgaattaaac aatcgtgttg gggaagcttt tcatgcatct tctactagtg tgttagtaga	1080
tggattttaca gatccttcaa ataacaaaag tagattctgc ctgggattgt tgtcaaatgt	1140
taatcgtaat tcaactattg aaaacactag gcggcatatt ggaaaagggtg ttcactctata	1200
ctacgttgggt ggggaggtgt acgctgagtg tcttagtgac agcagcatct ttgttcagag	1260
taggaactgc aactttcacc atggcttcca tcccaccacc gtctgtaaga tcccagcag	1320
ctgcagcctc aagattttta acaatcagga gtttgtcag cttctggctc agtcagtcaa	1380
ccatggattc gaggctgtgt atgagctcac caagatgtgt accattcgaa tgagctttgt	1440
caagggctgg ggagcagagt accaccgaca ggacgtcacc agtactccct gctggattga	1500
gattcacctc cacgggcctc tgcagtggct ggataaagtc cttactcaga tgggctctcc	1560
gctgaacccc atttcttctg tttcatagtg cagaagtatt ctttcaacta tatttttagt	1620
ggacttgttt taatttttaga ggaatttcca gtacagatgc tgtgagctga catggaaaac	1680
agatattatt tttctacgt aattgtgacc aacacatttg tattttatga tgatattaca	1740
tttgtttgta ttcgtgttca ttgtgattaa ctttcaaaag tattgtaaac gatgtagagt	1800
attttgcccc tgttgaaatg tttagcattg atcttaaact ggaacgtact ttttcttatt	1860
gtcccaacgt tttttaattt gttaaatttt ttttaciaaag tagttcatca cataatgaaa	1920
ttttatccta taagagaaca tatattgtgg aaagcagtag atgatatttc tctgggaatt	1980
tctttgcctt accacctttg aaaaagcata cattgtttgc aaaacctcaa agtagggctt	2040
gcttaaagga aactgttgaa tcttgtttga aggacactgc agtcctaacg tgttcagtga	2100
aagcaagggtg gtagatttct ggacgtcata catttacatt taatataggt aatattcatc	2160
agtgtaatgt gacttcatgc catatatatt ttgtaaaaca attccttttt aaaaacttca	2220
agtatttctc atttactcaa atttgttgta agtcctactt aacagttagt tactatgtgc	2280

tctgtggcct	tgttcagcat	tgtttgctgc	tttgggccaa	caattcaaga	actctaattt	2340
tcctgtgcat	taatcttttc	attttgcact	tttatgggtg	actgtcttag	tgtagcctct	2400
ggtaaaatac	tattaggtgg	cctggtttta	gagctcctcc	tcgctgcctt	ggcactcctt	2460
tgtgcaacac	gaccacttag	agatgacagc	tgtgagctgt	gctgcttttt	ctagccttta	2520
atttccaatg	tagtttataa	tgttgttctt	ctatagctcc	agctaagggtg	cctgttagtc	2580
ccctacaatg	ttatgagcat	tattgacatt	gaaaggttat	gtatgtatga	atacctttgc	2640
tccttaccag	acttgtcata	caaggactcg	tgcagtgtag	ccagtagagg	ctctttgggt	2700
ggcccaagaa	tgaggctggt	ggtgtaagtg	aatcacaata	gggattggga	tagttcatgt	2760
catatgtcat	atagcaagac	aatgtagagt	gtaggcttgt	ctctctgcat	caacgctctg	2820
cctctttctt	tttatccttt	tagaacctac	atggacgcta	atctccacaa	cactgttgga	2880
tgtgaacact	cttaagacac	tcatctagtt	cactgtgcct	tgtccttagg	actcttaacc	2940
actttctagg	gagcagttat	ggcctgagat	ggacagtcac	ggcctgagaa	tgaagacact	3000
actttgataa	agaaaaaggc	ctcatttgcc	tatcagagtgt	agaaagggtt	ttttctgggtg	3060
ccttttgaaa	atatacagag	ccacttgggt	cttctgctga	aaatgtaatt	ttggtttgac	3120
tttttagagt	gcccttcctg	cctttatgag	gaaaacagct	attttttttt	ttgggggggg	3180
ggattccttt	tgttttctgt	gccattattt	attgcctttc	agagtgaac	cattgggtgg	3240
ctttgctcct	tcagagaggg	ctccttgata	gccttcagta	gcttgagctg	tagacataag	3300
tattccatag	caagagtgtg	tcagctccat	gagagagatg	tctgctttat	agccgaggca	3360
gaaaccgttc	atgttccttt	acttggcagc	cttcaggaac	aggtttgtaa	gaacgtgtct	3420
tgagttgagt	gagtgtatgt	ctgtgagctc	tgtggaagtc	tggacacaag	ggccttgcc	3480
gctccttttt	tcagcagtg	gttacatgtt	gtctctccac	agtcttcatg	tcataggtct	3540
eggacttgca	gagtcctatg	tggcctgcc	tctgtacagt	ggcaggactg	aagctctgag	3600
ctgttctgag	gttcatggag	aaatcccaac	ctattctgtg	gtcagtaa	ggagactgtg	3660
tagtctacct	gctcctgtac	tgtccttact	gtatgtaagg	atatacagac	gcctgtgggt	3720
aggcagtact	cacagtgaga	tgaagacagc	aagtgtgcac	tgaaccacag	agggcaggg	3780
gtagggcctc	tgaagaagcc	accagaccag	accagtgccg	gtacagtctt	tgtcagagat	3840
ggctctgatg	gggccagac	tgaccctgac	catgctgagt	tgtgaggggt	agccttcagt	3900
tctctacct	ctgaagtgt	aggatgacag	acatccgcc	tcataccag	cttccgtgg	3960
gctaaggatc	agcctcagtc	ttcaggcgtg	ctaggcatgt	actttgccaa	gtatttagta	4020
tacaaaatac	attagtatct	gccagggaaa	aaagatttgc	aaataataaa	gattgccatc	4080

agtttgataa	atgttgtaaa	tggaagaatc	aaaatctcag	cgatggatta	cagcaacaag	4140
atgctgccta	ggaaaagcag	gaccaagagg	tacatttgac	tagtatacct	tcagcgtagc	4200
gtgatgacct	cactgatgtc	acccaactga	acttaagggc	tgtaagtagg	cgtgctgtgg	4260
gccttccaga	actagagaaa	attataggag	gaagtcagtt	ctaaagtatc	aaaagctggg	4320
taatggtggc	acatgccttt	gattctagca	ctcgggaagc	aggggctagc	ctagtctaca	4380
gagcaacttc	tacacagaga	aactgtcttg	gaggaaaata	aaaaaagaaa	agtcaaagag	4440
caaacaaata	gaacagagta	ggaatccgtg	tccccctttt	tctatgtttc	acggttgcag	4500
gtgtaagaaa	agtagtcata	gatgtggctg	agtttctaag	atgaaaccag	tagtaagatt	4560
gctaaatata	acacttcaac	caagttaaac	accctttggg	ggtatgaatg	aaagtaacac	4620
tgcaatatga	aatgaaccgt	gcaagtaaca	cttgggggta	cctcacagtc	tccctatgcc	4680
tgagaggact	gtgggaaaca	tttccatccc	ctgccagtat	cgccattggg	aggacagagt	4740
agatgaagaa	gtgaagtctt	actggtccag	ggcacgcctg	tcagcaatgc	catttgtgct	4800
tctgccacag	agagcaccga	gaggcttggc	tcagtatcct	cgaacccttct	ctggtcactt	4860
ccctggcagc	acttgggtcc	ctgtcactca	ctggtctctt	aaaagtcccg	tctctttgct	4920
tcctaaagat	tctctaaaaa	aattactatt	ttttatttct	tttttaaaag	tctttgttat	4980
tttggttttg	gatacagtct	ctttgtacag	tcctggctgg	cctggaaatt	actatgtagg	5040
ccagcctcaa	acttgaagta	attctcttgc	ctctgcctct	ggagttcttg	gattacaggc	5100
atgcactgca	gagtacagtg	agctctgatg	gcttttaaaa	ttcagcccct	ttgagggttt	5160
ggtttttagat	ccattagctt	tgtctgaacc	catctttgtc	cggccgagta	aatcctctgc	5220
tatccggggg	ctcggtagaa	atgtgttctc	agtatacata	cgactaaaca	ttggttgttt	5280
ataggtagcc	tcagatatatt	ggtagagcat	cttttttgaa	agtaatctcc	agctaggtgg	5340
gtatttccct	cacagcagta	ggattttccc	tttaggagat	accagttctt	catctttctt	5400
gtgaaaataa	tgcccttatg	gggagtgaag	attaaggagt	tgtttctaca	ctaacagaat	5460
tctatttgat	ggacaacttg	gacagtcttg	tggacttggg	tgggttctag	tgtgctaaga	5520
aggataacag	tattttaatag	tgtctgtcat	caggccttgc	tcattctccct	gtctagggct	5580
gtaggtcagt	gctcgagcac	ttagcaggca	tcgagtctag	tgttcagtgc	ccagcattgc	5640
acagaactca	gaatatatct	gtactgaaac	tgaagtgacc	acctacaacc	aggtggtatg	5700
ccagaaccac	agaaaggaga	ttcacgggtga	tgtgtttaaa	gcattgggct	ggtgacgggt	5760
gctgtgtagt	aatgacctct	tcctcagcaa	agagagtcct	ggagcaggct	gtcctcagaa	5820
gaggggaaggg	actggtgtgc	tccttgtgca	gataacttag	tgtataaatc	ggcatgagta	5880

```

gctatccttt aaggatttgt ttgaagttac tctttgtaaa aagttgagaa ttttgtgtgc 5940
agttgggcac atgcttgccc ttccccacc cgccatagtc ctgcctctct tgctgtgaac 6000
tggtgtcagc tacaacactc cagctaggtc tgagctcttt tgagagaagg tctcgtagag 6060
caccattctc agagagaagc taaagcatgg ggagccttag gacggtcagg caatgcactc 6120
tttaccacgg ctggctaagg ctgcagcttg accgtcctta cctaaatcag gtaagaatgt 6180
gattacagag cgagtgcttg tgttccccgg cctgccttct ccgaggaaga tgcttcatcc 6240
gaggatgatg cagagcagac gatggctgca ctgtaggctc gcctccttct gtgtatgggt 6300
tctgctgctg cttacggcat aggaaagta actagcagcg tgcttcaatt ctgccatctt 6360
ttgatactta taaaaatgta ttaggtttta ctgtattgtg ctctcaaagc cataactctt 6420
aagaaatttg gtttttttgc atattgtttg ctaatacttt gttttaataa acctcaaaat 6480
ctgcttaaaa aaaaaaaaaa aaaa 6504

```

```

<210> 21
<211> 465
<212> PRT
<213> Mus musculus

```

```

<400> 21

```

```

Met Thr Ser Met Ala Ser Leu Phe Ser Phe Thr Ser Pro Ala Val Lys
1          5          10          15

```

```

Arg Leu Leu Gly Trp Lys Gln Gly Asp Glu Glu Glu Lys Trp Ala Glu
          20          25          30

```

```

Lys Ala Val Asp Ala Leu Val Lys Lys Leu Lys Lys Lys Lys Gly Ala
          35          40          45

```

```

Met Glu Glu Leu Glu Lys Ala Leu Ser Ser Pro Gly Gln Pro Ser Lys
          50          55          60

```

```

Cys Val Thr Ile Pro Arg Ser Leu Asp Gly Arg Leu Gln Val Ser His
65          70          75          80

```

```

Arg Lys Gly Leu Pro His Val Ile Tyr Cys Arg Val Trp Arg Trp Pro
          85          90          95

```

```

Asp Leu Gln Ser His His Glu Leu Lys Pro Leu Asp Ile Cys Glu Phe
          100          105          110

```

```

Pro Phe Gly Ser Lys Gln Lys Glu Val Cys Ile Asn Pro Tyr His Tyr

```

115	120	125
Lys Arg Val Glu Ser Pro Val Leu Pro Pro Val Leu Val Pro Arg His 130 135 140		
Asn Glu Phe Asn Pro Gln His Ser Leu Leu Val Gln Phe Arg Asn Leu 145 150 155 160		
Ser His Asn Glu Pro His Met Pro Gln Asn Ala Thr Phe Pro Asp Ser 165 170 175		
Phe His Gln Pro Asn Asn Ala Pro Phe Pro Leu Ser Pro Asn Ser Pro 180 185 190		
Tyr Pro Pro Ser Pro Ala Ser Ser Thr Tyr Pro Asn Ser Pro Ala Ser 195 200 205		
Ser Gly Pro Gly Ser Pro Phe Gln Leu Pro Ala Asp Thr Pro Pro Pro 210 215 220		
Ala Tyr Met Pro Pro Asp Asp Gln Met Ala Pro Asp Asn Ser Gln Pro 225 230 235 240		
Met Asp Thr Ser Ser Asn Met Ile Pro Gln Thr Met Pro Ser Ile Ser 245 250 255		
Ser Arg Asp Val Gln Pro Val Ala Tyr Glu Glu Pro Lys His Trp Cys 260 265 270		
Ser Ile Val Tyr Tyr Glu Leu Asn Asn Arg Val Gly Glu Ala Phe His 275 280 285		
Ala Ser Ser Thr Ser Val Leu Val Asp Gly Phe Thr Asp Pro Ser Asn 290 295 300		
Asn Lys Ser Arg Phe Cys Leu Gly Leu Leu Ser Asn Val Asn Arg Asn 305 310 315 320		
Ser Thr Ile Glu Asn Thr Arg Arg His Ile Gly Lys Gly Val His Leu 325 330 335		
Tyr Tyr Val Gly Gly Glu Val Tyr Ala Glu Cys Leu Ser Asp Ser Ser 340 345 350		
Ile Phe Val Gln Ser Arg Asn Cys Asn Phe His His Gly Phe His Pro		

355 360 365
 Thr Thr Val Cys Lys Ile Pro Ser Ser Cys Ser Leu Lys Ile Phe Asn
 370 375 380
 Asn Gln Glu Phe Ala Gln Leu Leu Ala Gln Ser Val Asn His Gly Phe
 385 390 395 400
 Glu Ala Val Tyr Glu Leu Thr Lys Met Cys Thr Ile Arg Met Ser Phe
 405 410 415
 Val Lys Gly Trp Gly Ala Glu Tyr His Arg Gln Asp Val Thr Ser Thr
 420 425 430
 Pro Cys Trp Ile Glu Ile His Leu His Gly Pro Leu Gln Trp Leu Asp
 435 440 445
 Lys Val Leu Thr Gln Met Gly Ser Pro Leu Asn Pro Ile Ser Ser Val
 450 455 460
 Ser
 465

<210> 22
 <211> 2887
 <212> DNA
 <213> Homo sapiens

<400> 22
 gaattcggca cgaggagcga tcgagggagc tgagccgaga gaaagagccg ccgggcgctg 60
 cctcgccaga cctcgctggg accccggggc caccgggagg cacttttgtg gaggggggag 120
 gggggggcgac ctcggcagcc tcggcgcacg aagcgtccga gggcagcgtg ggggygggctg 180
 cgacctctgc atcggtggac tgcattttta attaaggatt ccagcagct ctttgggatt 240
 ttacagctt ccaactcatgt gttgacaccc gcgtccagga gaaactcgct ccaagtgcac 300
 ctagcgctg ggacctgaga cggcgttggc ctttcgtgca tgcaaatacca gggatttagg 360
 ttttgtttg gatttccttt tctttctttt cttttttttt tctttttgca gggagtaaga 420
 agggagctgg gggatatcaac aagcctgcct ttcggatcct gcgggaaaag cccatgtagt 480
 taagcgcttt ggttttaaaaa aaaggcaagg taaaggcagg gctttccaga cacatttagg 540
 ggttcgcgcg agctttgtgc tcatggacca gccgcacaac ttttgaaggc tcgccggccc 600
 atgtggggtc tttctggcgg gccgcctgca gccccctaa agcgcggggg gctggagttg 660
 ttgagcagcc ccgccgctgt ggtccatgta gccgctggcc gcgcgcggaa tgcggcttcg 720

cgtgcgcgtg	ttcccggccg	tcccgccttg	gcgaggtccc	tcatgttggt	gccctgcggc	780
gcccccttga	agacaggctg	tgcgcggttt	gcacggcggg	ccgcggcgga	ggttcatgtg	840
ggggtgcgac	ccgcgcagcc	ggcgccttgt	tgagggaacg	gacccccggg	aaccggagac	900
cgcctcccc	ccaccctgg	cgccaaagga	tattgtatgt	tcagggtccaa	acgctcgggg	960
ctggtgcggc	gactttggcg	aagtcgtgtg	gtccccaacc	gggaggaagg	cggcagcggc	1020
ggcggcgggtg	gcggcgacga	ggatgggagc	ttgggcagcc	gagctgagcc	ggccccgcgg	1080
gcaagagagg	gcggaggctg	cggccgctcc	gaagtccgcc	cggtagcccc	gcggcgggccc	1140
cgggacgcag	tgggacagcg	aggcgcccag	ggcgcgggga	ggcgccggcg	cgcagggggc	1200
cccccgaggc	ccatgtcgga	gccagggggc	ggcgtggga	gctccctgct	ggacgtggcg	1260
gagccgggag	gcccgggctg	gctgcccag	agtgactgcg	agacgggtgac	ctgctgtctc	1320
ttttcggagc	gggacgccgc	cggcgcgccc	cgggacgcca	gcgacccct	ggccggggcg	1380
gccctggagc	cggcgggcg	cggcgggagt	cgcgaagcgc	gctcgcggt	gctgctgctg	1440
gagcaggaac	tcaaaaccgt	cacgtactcg	ctgctgaagc	ggctcaagga	gcgctcgctg	1500
gacacgctgc	tggaggcggg	ggagtccgc	ggcggcgtgc	cgggcggctg	cgtgctggtg	1560
ccgcgcgccg	acctccgcct	gggcggccag	cccgcgccgc	cgcagctgct	gctcggccgc	1620
ctctttcgct	ggcccgacct	gcagcacgcc	gtggagctga	agccctgtg	cggctgccac	1680
agcttcgccg	ccgccgccga	cggccctacc	gtgtgctgca	acccctacca	cttcagccgg	1740
ctctgcgggc	ccgaatctcc	gccacctccc	tactctcggc	tgtctcctcg	cgacgagtac	1800
aagccactgg	atctgtccga	ttccacattg	tcttacactg	aaacggaggc	taccaactcc	1860
ctcatcactg	ctccgggtga	attctcagac	gccagcatgt	ctccggacgc	caccaagccg	1920
agccactggg	gcagcgtggc	gtactgggag	caccggacgc	gcgtgggccc	cctctatgcg	1980
gtgtacgacc	aggccgtcag	catcttctac	gacctacctc	agggcagcgg	cttctgcctg	2040
ggccagctca	acctggagca	gcgcagcgag	tcggtgcggc	gaacgcgcag	caagatcggc	2100
ttcggcatcc	tgtcagcaa	ggagcccag	ggcgtgtggg	cctacaaccg	cggcgagcac	2160
cccatcttcg	tcaactcccc	gacgtggac	gcgcccggcg	gccgcgccct	ggtcgtgcgc	2220
aaggtgcccc	ccggctactc	catcaagggt	ttcgacttcg	agcgctcggg	cctgcagcac	2280
gcgcccagac	ccgacgccgc	cgacggcccc	tacgacccca	acagcgtccg	catcagcttc	2340
gccaagggct	gggggccttg	ctactcccgg	cagttcatca	cctcctgccc	ctgctggctg	2400
gagatcctcc	tcaacaaccc	cagatagtgg	cggccccggc	gggagggggc	ggtgggaggc	2460
cgcgggccacc	gccacctgcc	ggcctcgaga	ggggccgatg	cccagagaca	cagccccccac	2520

```

ggacaaaacc cccagatat catctaccta gatttaatat aaagttttat atattatatg 2580
gaaatatata ttatacttgt aattatggag tcattttttac aatgtaatta tttatgtatg 2640
gtgcaatgtg tgtatatgga caaaacaaga aagacgcact ttggcttata attctttcaa 2700
tacagatata ttttctttct cttcctcctt cctcttcctt actttttata tatatatata 2760
aagaaaatga tacagcagag ctaggtggaa aagcctgggt ttggtgtatg gtttttgaga 2820
tattaatgcc cagacaaaaa gctaatacca gtcactcgat aataaagtat tcgcattata 2880
aaaaaga 2887

```

```

<210> 23
<211> 496
<212> PRT
<213> Homo sapiens

```

```

<400> 23

```

```

Met Phe Arg Ser Lys Arg Ser Gly Leu Val Arg Arg Leu Trp Arg Ser
1           5           10           15

```

```

Arg Val Val Pro Asn Arg Glu Glu Gly Gly Ser Gly Gly Gly Gly Gly
          20           25           30

```

```

Gly Asp Glu Asp Gly Ser Leu Gly Ser Arg Ala Glu Pro Ala Pro Arg
          35           40           45

```

```

Ala Arg Glu Gly Gly Gly Cys Gly Arg Ser Glu Val Arg Pro Val Ala
          50           55           60

```

```

Pro Arg Arg Pro Arg Asp Ala Val Gly Gln Arg Gly Ala Gln Gly Ala
65           70           75           80

```

```

Gly Arg Arg Arg Arg Ala Gly Gly Pro Pro Arg Pro Met Ser Glu Pro
          85           90           95

```

```

Gly Ala Gly Ala Gly Ser Ser Leu Leu Asp Val Ala Glu Pro Gly Gly
          100          105          110

```

```

Pro Gly Trp Leu Pro Glu Ser Asp Cys Glu Thr Val Thr Cys Cys Leu
          115          120          125

```

```

Phe Ser Glu Arg Asp Ala Ala Gly Ala Pro Arg Asp Ala Ser Asp Pro
          130          135          140

```

```

Leu Ala Gly Ala Ala Leu Glu Pro Ala Gly Gly Gly Arg Ser Arg Glu

```


145 150 155 160
 Ala Arg Ser Arg Leu Leu Leu Leu Glu Gln Glu Leu Lys Thr Val Thr
 165 170 175
 Tyr Ser Leu Leu Lys Arg Leu Lys Glu Arg Ser Leu Asp Thr Leu Leu
 180 185 190
 Glu Ala Val Glu Ser Arg Gly Gly Val Pro Gly Gly Cys Val Leu Val
 195 200 205
 Pro Arg Ala Asp Leu Arg Leu Gly Gly Gln Pro Ala Pro Pro Gln Leu
 210 215 220
 Leu Leu Gly Arg Leu Phe Arg Trp Pro Asp Leu Gln His Ala Val Glu
 225 230 235 240
 Leu Lys Pro Leu Cys Gly Cys His Ser Phe Ala Ala Ala Ala Asp Gly
 245 250 255
 Pro Thr Val Cys Cys Asn Pro Tyr His Phe Ser Arg Leu Cys Gly Pro
 260 265 270
 Glu Ser Pro Pro Pro Pro Tyr Ser Arg Leu Ser Pro Arg Asp Glu Tyr
 275 280 285
 Lys Pro Leu Asp Leu Ser Asp Ser Thr Leu Ser Tyr Thr Glu Thr Glu
 290 295 300
 Ala Thr Asn Ser Leu Ile Thr Ala Pro Gly Glu Phe Ser Asp Ala Ser
 305 310 315 320
 Met Ser Pro Asp Ala Thr Lys Pro Ser His Trp Cys Ser Val Ala Tyr
 325 330 335
 Trp Glu His Arg Thr Arg Val Gly Arg Leu Tyr Ala Val Tyr Asp Gln
 340 345 350
 Ala Val Ser Ile Phe Tyr Asp Leu Pro Gln Gly Ser Gly Phe Cys Leu
 355 360 365
 Gly Gln Leu Asn Leu Glu Gln Arg Ser Glu Ser Val Arg Arg Thr Arg
 370 375 380
 Ser Lys Ile Gly Phe Gly Ile Leu Leu Ser Lys Glu Pro Asp Gly Val

385 390 395 400
 Trp Ala Tyr Asn Arg Gly Glu His Pro Ile Phe Val Asn Ser Pro Thr
 405 410 415
 Leu Asp Ala Pro Gly Gly Arg Ala Leu Val Val Arg Lys Val Pro Pro
 420 425 430
 Gly Tyr Ser Ile Lys Val Phe Asp Phe Glu Arg Ser Gly Leu Gln His
 435 440 445
 Ala Pro Glu Pro Asp Ala Ala Asp Gly Pro Tyr Asp Pro Asn Ser Val
 450 455 460
 Arg Ile Ser Phe Ala Lys Gly Trp Gly Pro Cys Tyr Ser Arg Gln Phe
 465 470 475 480
 Ile Thr Ser Cys Pro Cys Trp Leu Glu Ile Leu Leu Asn Asn Pro Arg
 485 490 495

<210> 24
 <211> 3103
 <212> DNA
 <213> Homo sapiens

<400> 24
 cggagagccg cgcagggcgc gggccgcgcg ggggtggggca gccggagcgc agggccccga 60
 tccccggcgg gcgcccccg gccccgcgc gcgccccggc ctccgggaga ctggcgcatg 120
 ccacggagcg cccctcgggc cgcgcgcgct cctgccccgg cccctgctgc tgctgctgtc 180
 gcctgcgcct gctgccccaa ctcggcgccc gactttcttca tgggtgtgcgg aggtcatggt 240
 cgctccttag caggcaaacg acttttctcc tcgcctctc gcccgcatg ttcaggacca 300
 aacgatctgc gctcgtccgg cgtctctgga ggagccgtgc gcccggcggc gaggacgagg 360
 aggagggcgc agggggagggt ggaggaggag gcgagctgcg gggagaagg gcgacggaca 420
 gccgagcgca tggggccgggt ggcggcgggc cgggcagggc tggatgctgc ctgggcaagg 480
 cgggtgcgagg tgccaaagg caccaccatc cccaccgcc agccgcgggc gccggcgcg 540
 ccggggggcgc cgaggcgat ctgaaggcgc tcacgcactc ggtgctcaag aaactgaagg 600
 agcggcagct ggagctgctg ctccaggccg tggagtcccc cggcgggacg cgcaccgcgt 660
 gcctcctgct gcccggccgc ctggactgca ggctggggcc gggggcgccc gccggcgcg 720
 agcctgcgca gccgcctcgc tctactcgc tccccctcct gctgtgcaaa gtgttcagg 780
 ggccggatct caggcattcc tcggaagtca agaggctgtg ttgctgtgaa tcttacggga 840

```

agatcaaccc cgagctggtg tgctgcaacc cccatcacct tagccgactc tgcgaactag      900
agtctccccc cctccttac tccagatacc cgatggattt tctcaaacca actgcagact      960
gtccagatgc tgtgccttc tccgctgaaa cagggggaac gaattatctg gcccctgggg      1020
ggctttcaga ttcccaactt cttctggagc ctggggatcg gtcacactgg tgcgtggtgg      1080
catactggga ggagaagacg agagtgggga ggctctactg tgtccaggag ccctctctgg      1140
atatcttcta tgatctacct caggggaatg gcttttgctt cggacagctc aattcggaca      1200
acaagagtca gctggtgcag aaggtgcgga gcaaaatcgg ctgcggcatc cagctgacgc      1260
gggaggtgga tgggtgtgtg gtgtacaacc gcagcagtta ccccatcttc atcaagtccg      1320
ccacactgga caaccggac tccaggacgc tgttgggtaca caaggtgttc cccggtttct      1380
ccatcaaggc tttcgactac gagaaggcgt acagcctgca gcggcccaat gaccacgagt      1440
ttatgcagca gccgtggacg ggctttaccg tgcagatcag ctttgtgaag ggctggggtc      1500
agtgtctacac ccgccagttc atcagcagct gcccgctgtg gctagaggtc atcttcaaca      1560
gccggtagcc gcgtgcggag gggacagagc gtgagctgag caggccacac ttcaaactac      1620
tttgctgcta atattttctt cctgagtgtt tgcttttcat gcaaactctt tggtcgtttt      1680
ttttttgttt gttggttggg tttcttcttc tgcctctcgt ttgtgttctg ttttgtttcg      1740
ctctttgaga aatagcttat gaaaagaatt gttggggggt tttttggaag aaggggcagg      1800
tatgatcggc aggacacct gataggaaga ggggaagcag aaatccaagc accaccaaac      1860
acagtgtatg aagggggggc gtcattctt cacttgtcag gagtgtgtgt gagtgtgagt      1920
gtgcggctgt gtgtgcacgc gtgtgcagga gcggcagatg gggagacaac gtgctctttg      1980
ttttgtgtct cttatggatg tcccagcag agaggtttgc agtcccaagc ggtgtctctc      2040
ctgccccttg gacacgctca gtggggcaga ggcagtacct gggcaagctg gcggctgggg      2100
tcccagcagc tgccaggagc acggctctgt cccagcctg ggaaagcccc tgcccctcct      2160
ctccctcatc aaggacacgg gcctgtccac aggtctctga gcagcgagcc tgctagtggc      2220
cgaaccagaa ccaattatth tcatccttgt cttattccct tctgccagc ccctgccatt      2280
gtagcgtctt tcttttttgg ccatctgtct ctggatctcc ctgagatggg cttcccaagg      2340
gctgccgggg cagccccctc acagtattgc tcaccagtg ccctctcccc tcagcctctc      2400
ccctgcctgc cctggtgaca tcagggtttt cccggactta gaaaaccagc tcagcactgc      2460
ctgctcccat cctgtgtgtt aagctctgct attaggccag caagcgggga tgtccctggg      2520
agggacatgc ttagcagtc ccttccctcc aagaaggatt tggtcctgca taaccaagg      2580
taccatccta ggctgacacc taactcttct ttcatttctt ctacaactca tacactcgta      2640

```

```

tgataacttcg acactgttct tagctcaatg agcatgttta gactttaaca taagctatct 2700
ttctaactac aaaggtttaa atgaacaaga gaagcattct cattggaaat ttagcattgt 2760
agtgccttga gagagaaagg actcctgaaa aaaaacctga gatttattaa agaaaaaaat 2820
gtatatttatg ttatatataa atatattatt acttgtaaata ataaagacgt tttataagca 2880
tcattatttta tgtattgtgc aatgtgtata aacaagaaaa ataaagaaaa gatgcacttt 2940
gctttaatat aaatgcaaata aacaaatgcc aaattaaaaa agataaacac aagattgggtg 3000
ttttttccta tgggtgttat cacctagctg aatgtttttc taaaggagtt tatgttccat 3060
taaacgattt ttaaaatgta cacttgaaaa aaaaaaaaaa aaa 3103

```

```

<210> 25
<211> 426
<212> PRT
<213> Homo sapiens

```

```

<400> 25

```

```

Met Phe Arg Thr Lys Arg Ser Ala Leu Val Arg Arg Leu Trp Arg Ser
1           5           10           15

```

```

Arg Ala Pro Gly Gly Glu Asp Glu Glu Glu Gly Ala Gly Gly Gly Gly
          20           25           30

```

```

Gly Gly Gly Glu Leu Arg Gly Glu Gly Ala Thr Asp Ser Arg Ala His
35           40           45

```

```

Gly Ala Gly Gly Gly Gly Pro Gly Arg Ala Gly Cys Cys Leu Gly Lys
50           55           60

```

```

Ala Val Arg Gly Ala Lys Gly His His His Pro His Pro Pro Ala Ala
65           70           75           80

```

```

Gly Ala Gly Ala Ala Gly Gly Ala Glu Ala Asp Leu Lys Ala Leu Thr
          85           90           95

```

```

His Ser Val Leu Lys Lys Leu Lys Glu Arg Gln Leu Glu Leu Leu Leu
100           105           110

```

```

Gln Ala Val Glu Ser Arg Gly Gly Thr Arg Thr Ala Cys Leu Leu Leu
115           120           125

```

```

Pro Gly Arg Leu Asp Cys Arg Leu Gly Pro Gly Ala Pro Ala Gly Ala
130           135           140

```

Gln Pro Ala Gln Pro Pro Ser Ser Tyr Ser Leu Pro Leu Leu Leu Cys
 145 150 155 160
 Lys Val Phe Arg Trp Pro Asp Leu Arg His Ser Ser Glu Val Lys Arg
 165 170 175
 Leu Cys Cys Cys Glu Ser Tyr Gly Lys Ile Asn Pro Glu Leu Val Cys
 180 185 190
 Cys Asn Pro His His Leu Ser Arg Leu Cys Glu Leu Glu Ser Pro Pro
 195 200 205
 Pro Pro Tyr Ser Arg Tyr Pro Met Asp Phe Leu Lys Pro Thr Ala Asp
 210 215 220
 Cys Pro Asp Ala Val Pro Ser Ser Ala Glu Thr Gly Gly Thr Asn Tyr
 225 230 235 240
 Leu Ala Pro Gly Gly Leu Ser Asp Ser Gln Leu Leu Leu Glu Pro Gly
 245 250 255
 Asp Arg Ser His Trp Cys Val Val Ala Tyr Trp Glu Glu Lys Thr Arg
 260 265 270
 Val Gly Arg Leu Tyr Cys Val Gln Glu Pro Ser Leu Asp Ile Phe Tyr
 275 280 285
 Asp Leu Pro Gln Gly Asn Gly Phe Cys Leu Gly Gln Leu Asn Ser Asp
 290 295 300
 Asn Lys Ser Gln Leu Val Gln Lys Val Arg Ser Lys Ile Gly Cys Gly
 305 310 315 320
 Ile Gln Leu Thr Arg Glu Val Asp Gly Val Trp Val Tyr Asn Arg Ser
 325 330 335
 Ser Tyr Pro Ile Phe Ile Lys Ser Ala Thr Leu Asp Asn Pro Asp Ser
 340 345 350
 Arg Thr Leu Leu Val His Lys Val Phe Pro Gly Phe Ser Ile Lys Ala
 355 360 365
 Phe Asp Tyr Glu Lys Ala Tyr Ser Leu Gln Arg Pro Asn Asp His Glu
 370 375 380

Phe Met Gln Gln Pro Trp Thr Gly Phe Thr Val Gln Ile Ser Phe Val
 385 390 395 400

Lys Gly Trp Gly Gln Cys Tyr Thr Arg Gln Phe Ile Ser Ser Cys Pro
 405 410 415

Cys Trp Leu Glu Val Ile Phe Asn Ser Arg
 420 425

<210> 26
 <211> 1600
 <212> DNA
 <213> Mus musculus

<400> 26
 tttgaatcgg accccactgc ttccagaagg aaaggcaagc ttgttcctat gacatccgtg 60
 gacaggtact tgccgccgac ctgcccgagg ccctgcaagc cttgaaaggt ctcatcctct 120
 ttccccgtgc agcagcctga gctctgcctc ctatgcaccc cagcaccccc atcagctccc 180
 tcttctcctt caccagcccc gcagtgaagc ggctgctggg ctggaagcag ggagatgaag 240
 aggagaagtg ggcagagaag gcggtggact ctttggtgaa gaagttaaag aagaagaaag 300
 gcgccatgga tgaactggag agggcgctga gctgcccggg tcagcctagc aagtgtgtca 360
 ccatcccaag gtccctcgat ggacgcctcc aggtgtccca ccgaaagggg ctgccccacg 420
 tcatctactg ccgcgtgtgg cgctggccag acctgcagtc ccatcatgag ctgaagccct 480
 tggagtgtcg tgagttcccc ttccggctcca agcagaagga ggtctgcac aaccataacc 540
 attaccgcag agtggagacc ccagttctgc ctccagtgtc ggtaccaaga cacagcgagt 600
 acaaccctca gctcagcctc ctggccaagt tccgaagtgc ctgcgtgcac agcgaacccc 660
 tcatgccgca caacgccacc taccctgact ctttccagca gtctctctgt ccggcaccgc 720
 cctcctcgcc aggccatgtg tttccgcagt ctccatgccc caccagctac ccgcaactccc 780
 ccggaagtcc ttccgagtca gacagtcctt atcaacactc agacttccgg ccagtttgtc 840
 acgaggaacc ccagcactgg tgttctgttg cctactacga actaaacaac cgggtcggag 900
 agactttcca ggcgtcctcg cggagcgtgc tcatagacgg cttcaccgac ctttccaata 960
 acaggaatag gttttgcctt gggcttctct caaatgtaaa cagaaactcg accatagaaa 1020
 acaccaggag gcacattgga aaggggtgtgc atttgtacta cgttgggggc gaggtgtatg 1080
 cggagtgcgt gagcgacagc agcatctttg tccagagccg gaactgcaac taccagcacg 1140
 gcttccaccc ggccaccgtc tgcaagatcc ccagcggctg cagcctcaag gtcttcaaca 1200
 accagctctt cgcgccagctg ctcgcccagt ccgtgcacca cggctttgaa gtggtgtatg 1260

```

agctgacgaa gatgtgcacg attcggatga gctttgtgaa gggctgggga gcagagtatc 1320
atcgccagga tgtcacgagc accccctgct ggatcgagat ccatcttcat ggaccgctgc 1380
agtggttgga taagggtgctc actcagatgg gctccccaca caaccctatc tcttcagtgt 1440
cttaagtcac gtcgtcagcc acgttgccac agaacagact cgggcagggg cttccatcgt 1500
ggcaaccgca gctaatagcag ggttccggat gcagatgtaa atacacgtgt aacgcatccg 1560
agtcacgttt atatcaccgt ttgttttgtg ctacctactt 1600

```

<210> 27

<211> 430

<212> PRT

<213> Mus musculus

<400> 27

```

Met His Pro Ser Thr Pro Ile Ser Ser Leu Phe Ser Phe Thr Ser Pro
1          5          10          15

```

```

Ala Val Lys Arg Leu Leu Gly Trp Lys Gln Gly Asp Glu Glu Glu Lys
20          25          30

```

```

Trp Ala Glu Lys Ala Val Asp Ser Leu Val Lys Lys Leu Lys Lys Lys
35          40          45

```

```

Lys Gly Ala Met Asp Glu Leu Glu Arg Ala Leu Ser Cys Pro Gly Gln
50          55          60

```

```

Pro Ser Lys Cys Val Thr Ile Pro Arg Ser Leu Asp Gly Arg Leu Gln
65          70          75          80

```

```

Val Ser His Arg Lys Gly Leu Pro His Val Ile Tyr Cys Arg Val Trp
85          90          95

```

```

Arg Trp Pro Asp Leu Gln Ser His His Glu Leu Lys Pro Leu Glu Cys
100         105         110

```

```

Cys Glu Phe Pro Phe Gly Ser Lys Gln Lys Glu Val Cys Ile Asn Pro
115         120         125

```

```

Tyr His Tyr Arg Arg Val Glu Thr Pro Val Leu Pro Pro Val Leu Val
130         135         140

```

```

Pro Arg His Ser Glu Tyr Asn Pro Gln Leu Ser Leu Leu Ala Lys Phe
145         150         155         160

```

Arg Ser Ala Ser Leu His Ser Glu Pro Leu Met Pro His Asn Ala Thr
 165 170 175
 Tyr Pro Asp Ser Phe Gln Gln Ser Leu Cys Pro Ala Pro Pro Ser Ser
 180 185 190
 Pro Gly His Val Phe Pro Gln Ser Pro Cys Pro Thr Ser Tyr Pro His
 195 200 205
 Ser Pro Gly Ser Pro Ser Glu Ser Asp Ser Pro Tyr Gln His Ser Asp
 210 215 220
 Phe Arg Pro Val Cys Tyr Glu Glu Pro Gln His Trp Cys Ser Val Ala
 225 230 235 240
 Tyr Tyr Glu Leu Asn Asn Arg Val Gly Glu Thr Phe Gln Ala Ser Ser
 245 250 255
 Arg Ser Val Leu Ile Asp Gly Phe Thr Asp Pro Ser Asn Asn Arg Asn
 260 265 270
 Arg Phe Cys Leu Gly Leu Leu Ser Asn Val Asn Arg Asn Ser Thr Ile
 275 280 285
 Glu Asn Thr Arg Arg His Ile Gly Lys Gly Val His Leu Tyr Tyr Val
 290 295 300
 Gly Gly Glu Val Tyr Ala Glu Cys Val Ser Asp Ser Ser Ile Phe Val
 305 310 315 320
 Gln Ser Arg Asn Cys Asn Tyr Gln His Gly Phe His Pro Ala Thr Val
 325 330 335
 Cys Lys Ile Pro Ser Gly Cys Ser Leu Lys Val Phe Asn Asn Gln Leu
 340 345 350
 Phe Ala Gln Leu Leu Ala Gln Ser Val His His Gly Phe Glu Val Val
 355 360 365
 Tyr Glu Leu Thr Lys Met Cys Thr Ile Arg Met Ser Phe Val Lys Gly
 370 375 380
 Trp Gly Ala Glu Tyr His Arg Gln Asp Val Thr Ser Thr Pro Cys Trp
 385 390 395 400

Ile Glu Ile His Leu His Gly Pro Leu Gln Trp Leu Asp Lys Val Leu
405 410 415

Thr Gln Met Gly Ser Pro His Asn Pro Ile Ser Ser Val Ser
420 425 430

INTERNATIONAL SEARCH REPORT

In Application No
PCT/US2005/003229

A. CLASSIFICATION OF SUBJECT MATTER
IPC 7 A61K38/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	TAMAKI ET AL: "Intracellular signaling of osteogenic protein-1 through Smad5 activation" JOURNAL OF CELLULAR PHYSIOLOGY, vol. 177, 1998, pages 355-363, XP008048090 * See page 355 (Abstract) *	1-5,7-49
Y	AOKI ET AL: "Synergistic effects of different bone morphogenetic protein type I receptors on alkaline phosphatase induction" JOURNAL OF CELL SCIENCE, vol. 114, 2001, pages 1483-1489, XP002302367 * See pages 1483-1484 (Abstract and Introduction, esp. last sentence), figures 1-5, and pages 1487-1488 (Discussion) *	1-5,7-49

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

° Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- "&" document member of the same patent family

Date of the actual completion of the international search

7 June 2005

Date of mailing of the international search report

17/06/2005

Name and mailing address of the ISA
European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Korsner, S-E

INTERNATIONAL SEARCH REPORT

Int I Application No
PCT/US2005/003229

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	<p>YEH ET AL: "Osteogenic protein-1 (OP-1, BMP-7) induces osteoblastic cell differentiation of the pluripotent mesenchymal cell line C2C12"</p> <p>JOURNAL OF CELLULAR BIOCHEMISTRY, vol. 87, 2002, pages 292-304, XP002302368</p> <p>* See pages 300-303 (Discussion) *</p>	1-5,7-49
Y	<p>FORSLUND ET AL: "A comparative dose-response study of cartilage-derived morphogenetic protein (CDMP)-1, -2 and -3 for tendon healing in rats"</p> <p>JOURNAL OF ORTHOPAEDIC RESEARCH, vol. 21, 2003, pages 617-621, XP002330748</p> <p>* See pages 617-618 (Introduction), and pages 619-621 (Discussion) *</p>	1-5,7-49
Y	<p>TAKASE ET AL: "Induction of Smad6 mRNA by bone morphogenetic proteins"</p> <p>BIOCHEMICAL AND BIOPHYSICAL RESEARCH COMMUNICATIONS, vol. 244, 1998, pages 26-29, XP002219536</p> <p>* See page 26 (Abstract) *</p>	1-5,7-49
Y	<p>MACÍAS-SILVA ET AL: "Specific activation of Smad1 signaling pathways by the BMP7 type I receptor, ALK2"</p> <p>THE JOURNAL OF BIOLOGICAL CHEMISTRY, vol. 273, 1998, pages 25628-25636, XP002330481</p> <p>* See page 25630 (right column) - page 25631 (left column) *</p>	1-5,7-49
Y	<p>NISHIDA ET AL: "Osteogenic protein 1 stimulates cell-associated matrix assembly by normal human articular chondrocytes"</p> <p>ARTHRITIS & RHEUMATISM, vol. 43, 2000, pages 206-214, XP002330479</p> <p>* See pages 206-207 (Introduction) *</p>	1-5,7-49
Y	<p>SIMANK ET AL: "Das Wachstumsfaktorkomposit aus GDF-5 und mineralisiertem Kollagen verbessert die Ausheilung einer Hüftkopfnekrose"</p> <p>ORTHOPÄDE, vol. 33, January 2004 (2004-01), pages 68-75, XP002330749</p> <p>* See page 71 (English summary) *</p>	1-5,7-49
Y	<p>US 6 531 445 B1 (COHEN ET AL) 11 March 2003 (2003-03-11)</p> <p>* See columns 2-3 (tissues) and 6-7 (proteins useful) *</p>	1-5,7-49
	-/--	

INTERNATIONAL SEARCH REPORT

Inte

I Application No

PCT/US2005/003229

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 03/059932 A (YISSUM RESEARCH DEV. COMP. OF THE HEBREW UNIV. / GBF-GESELLSCHAFT) 24 July 2003 (2003-07-24) * See page 1 (Field of the invention), pages 11 and 21 *	1-5,7-49
P,X	YEH ET AL: "Cartilage-derived morphogenetic proteins enhance the osteogenic protein-1-induced osteoblastic cell differentiation of C2C12 cells" JOURNAL OF CELLULAR PHYSIOLOGY, vol. 201, December 2004 (2004-12), pages 401-408, XP008048134 * See page 401 (Abstract) *	1-5,7-49

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US2005/003229

Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

Although Claims 1-49 (partially) are directed
to a method of treatment of the human body,
the search has been carried out and based
on the alleged effects of the compounds.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such
an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all
searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment
of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report
covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is
restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Inter

Application No

PCT/US2005/003229

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 6531445	B1	11-03-2003	US 2004102373 A1 27-05-2004
			US 6565843 B1 20-05-2003
			AT 192931 T 15-06-2000
			AU 669127 B2 30-05-1996
			AU 2564592 A 05-04-1993
			AU 670558 B2 25-07-1996
			AU 3176293 A 27-04-1993
			CA 2104678 A1 12-09-1992
			CA 2116559 A1 01-04-1993
			CA 2116562 A1 18-03-1993
			CA 2363965 A1 17-09-1992
			DE 69231062 D1 21-06-2000
			DE 69231062 T2 15-02-2001
			EP 0601106 A1 15-06-1994
			EP 0601135 A1 15-06-1994
			ES 2149776 T3 16-11-2000
			JP 6510989 T 08-12-1994
			JP 7502021 T 02-03-1995
			US 2003125230 A1 03-07-2003
			WO 9304692 A1 18-03-1993
			WO 9305751 A2 01-04-1993
			US 2003105004 A1 05-06-2003
			US 5656593 A 12-08-1997
			US 6333312 B1 25-12-2001
			US 6153583 A 28-11-2000
			US 2003104993 A1 05-06-2003
			US 6211146 B1 03-04-2001
			US 2003109445 A1 12-06-2003
			US 5650276 A 22-07-1997
			US 6022853 A 08-02-2000
			US 6395883 B1 28-05-2002
			US 5674844 A 07-10-1997
			US 6077823 A 20-06-2000
			US 5849686 A 15-12-1998
			US 5741641 A 21-04-1998
			US 6495513 B1 17-12-2002
			US 5834179 A 10-11-1998
			US 6399569 B1 04-06-2002
			US 5972884 A 26-10-1999
			US 5739107 A 14-04-1998
			US 6090776 A 18-07-2000
			US 5733878 A 31-03-1998
			US 2005065083 A1 24-03-2005
			US 2005096339 A1 05-05-2005
			US 6071708 A 06-06-2000
			US 5854071 A 29-12-1998
			US 5994131 A 30-11-1999
			US 2002049159 A1 25-04-2002
			US 6506729 B1 14-01-2003
WO 03059932	A	24-07-2003	AU 2002366974 A1 30-07-2003
			CA 2471891 A1 24-07-2003
			EP 1467624 A2 20-10-2004
			WO 03059932 A2 24-07-2003
			US 2003228292 A1 11-12-2003
			US 2003219423 A1 27-11-2003